

August 16, 1996 / Vol. 45 / No. 32

- 689 Adverse Events Associated with Ephedrine-Containing Products — Texas
- 693 Update: Diphtheria Epidemic New Independent States of the Former Soviet Union, January 1995– March 1996
- 697 Progress Toward Poliomyelitis Eradication — Indonesia, 1995
- 700 Average Postpartum Length of Stay for Uncomplicated Deliveries — New Jersey, 1995

Adverse Events Associated with Ephedrine-Containing Products — Texas, December 1993–September 1995

MORBIDITY AND MORTALITY WEEKLY REPORT

During December 1993–September 1995, the Bureau of Food and Drug Safety, Texas Department of Health (TDH), received approximately 500 reports of adverse events in persons who consumed dietary supplement products containing ephedrine and associated alkaloids (pseudoephedrine, norephedrine, and N-methyl ephedrine). This total included reports by individuals and reports identified by the Bureau of Epidemiology, TDH, in a review of records from the six centers of the Texas Poison Center Network. Reported adverse events ranged in severity from tremor and headache to death in eight ephedrine users and included reports of stroke, myocardial infarction, chest pain, seizures, insomnia, nausea and vomiting, fatigue, and dizziness. Seven of the eight reported fatalities were attributed to myocardial infarction or cerebrovascular accident. This report describes three patients in which the recommended dosage for the dietary supplements reportedly was not exceeded, summarizes results from ongoing investigations, and underscores the potential health risks associated with the use of products containing ephedrine.

Case Reports

Patient 1. In December 1993, a 44-year-old man died from acute coronary artery thrombosis approximately 3 weeks after beginning daily use of a dietary supplement containing ephedrine. He was an active swimmer and tennis player with no known cardiovascular risk factors. He received the dietary supplement from his family physician during a routine physical examination when he requested a substitute for his daily coffee and cocoa. He used the product as directed and eliminated his coffee and cocoa use. On December 18, 1993, after playing tennis and returning home, he sustained a cardiorespiratory arrest. An autopsy revealed an acute thrombus in the left anterior descending coronary artery. All other coronary lumina were patent, although calcified with focal narrowing to approximately 50%.

Patient 2. In May 1995, a 35-year-old woman who was taking no other prescription or over-the-counter (OTC) medications began use of a dietary supplement containing ephedrine for weight loss. She used the supplement within the dosage recommended on the label for approximately 30 days, discontinued use of the supplement while on a 1-week vacation, then resumed the usual dosage when she returned on June 24, 1995. On June 25, while sleeping, she had acute onset of symptoms including anterior

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES / Public Health Service

Ephedrine — Continued

chest pain that radiated to her left shoulder and arm, numbness of the left arm and hand, diaphoresis, and shortness of breath. She was taken to the hospital and her pain remitted after she was treated with nitroglycerin and morphine. Although an electrocardiogram and cardiac enzymes indicated an acute myocardial infarct, cardiac catheterization indicated normal cardiac function and normal coronary arteries. She had no history of cardiovascular risk factors. She was discharged with a diagnosis of acute myocardial infarction secondary to cardiac spasm and was advised to discontinue use of the dietary supplement that contained ephedrine. Since discontinuing use of the product, she has had no additional cardiac-related symptoms.

Patient 3. On August 17, 1995, a 38-year-old woman with no history of seizures experienced two petit mal seizures beginning at 11 p.m. She experienced two additional petit mal seizures the following morning, and that afternoon had onset of a generalized tonic-clonic seizure lasting approximately 2 minutes, during which she required respiratory assistance. On August 17, she had taken two tablets of an ephedrine-containing dietary supplement at 10 a.m. and two more 5 hours later as directed on the product label. She denied use of other drugs except oral contraceptives. During August 19–22, she experienced five additional episodes of unresponsiveness while sitting or standing; while waiting in the office of a neurologist, she sustained an additional generalized seizure witnessed by the neurologist and staff. She was hospitalized for monitoring, treated with antiseizure medication, and diagnosed with new onset of tonic-clonic seizures with complex partial seizures. Other possible causes of seizures were excluded. She was discharged and was advised to avoid any medications or products that contained ephedrine, pseudoephedrine, or related drugs. Since discontinuing use of the product, she has had no additional seizures.

Ongoing Investigations

TDH also has received reports of persons who had acute onset of palpitations and fainting after using ephedrine-containing products marketed as "beyond smart drugs" for "euphoric stimulation, highly increased energy levels, tingly skin sensations, enhanced sensory processing, increased sexual sensations, and mood elevations." Although these substances have been sold without warnings or contraindications on the information labels, one label indicated that the product "acts on the same basis as MDMA (3,4-methylenedioxy-methamphetamine, "ecstasy") triggering similar but not identical physical reactions in the body." TDH investigators purchased a product labeled "no side effects" that also listed wild Chinese ginseng as the only ingredient. Laboratory analysis indicated that a single tablet contained 45 mg ephedrine and 20 mg caffeine; the label on this product instructed users to take five tablets, representing a total ephedrine dosage of approximately 11 times the usual recommended OTC dosage of bronchodilator products, which contain 12.5 mg–25.0 mg of ephedrine per dose.

Ephedrine-containing products usually are marketed and labeled for weight loss, energy, "pep," performance enhancement, or body building or as a substitute for illicit drugs such as MDMA. They are commonly labeled as "natural" or "herbal" and use common names for herbs as the source of active ingredients (ma huang, Chinese ephedra, and *Sida cordifolia*—another plant source with small amounts of ephedrine alkaloids). An additional 400 reports of adverse events involved OTC drug products

Ephedrine — Continued

containing ephedrine that were labeled as required for use as bronchodilators but marketed in a manner to imply their effectiveness for weight loss and as stimulants.

Since September 1995, the Texas Poison Control Network has received approximately 300 additional reports of adverse events in persons consuming products containing ephedrine. These reports are being investigated by TDH.

Reported by: DM Perrotta, PhD, Bur of Epidemiology; G Coody, C Culmo, Bur of Food and Drug Safety; Texas Poison Center Network, Texas Dept of Health. Clinical Research and Review Staff, Center for Food Safety and Applied Nutrition, Food and Drug Administration. Environmental Hazards Epidemiology Section, Health Studies Br, Div of Environmental Hazards and Health Effects, National Center for Environmental Health, CDC.

Editorial Note: The three reports presented here and the approximately 500 reports of adverse events received by TDH underscore that use of dietary supplements containing ephedrine and related alkaloids can be associated with a spectrum of adverse health events. Although a cause-and-effect relation cannot be established for the three patients presented here, no other cause was found to explain their medical conditions, all of which are compatible with documented effects of ephedrine consumption. Ephedrine and associated alkaloids are structurally similar to the amphetamines (1) and, by stimulating adrenergic receptors, can increase arterial blood pressure through both peripheral vasoconstriction and cardiac stimulation. Adverse effects from ephedrine can be variable, and do not always depend on the dose consumed. Serious adverse effects of ephedrine and related alkaloids, such as acute cardiovascular and central nervous system stimulant effects, can occur in susceptible persons with use of low dosages. Other adverse effects associated with the use of ephedrine include palpitations, tachycardia, hypertension, coronary spasm, paranoid psychoses, convulsions, respiratory depression, coma, and death (2). Particularly when used in combinations with phenylpropanolamine (PPA) and caffeine, ephedrine has been associated with stroke secondary to intracranial hemorrhage, seizures, mania, and psychosis (3,4). Combinations of ephedrine and caffeine have been documented to have side effects substantially greater than those from the consumption of either compound alone or of a placebo (5-8).

In the United States, ephedrine, pseudoephedrine, and PPA have been marketed extensively for some OTC uses. For example, preparations containing ephedrine are marketed for oral use as a short-term, OTC bronchodilator for persons with mild asthma. The Food and Drug Administration (FDA) has proposed to remove oral ephedrine drug products from the OTC market based on their use in the production of illicit drugs and on their misuse and abuse as stimulants and for weight loss.* Pseudoephedrine, an ephedrine alkaloid contained in many OTC decongestant, cold, and allergy products, is associated with fewer cardiovascular and central nervous system stimulant effects than ephedrine. PPA, another ephedrine alkaloid, also is contained in OTC decongestant, cold, and allergy preparations and is marketed for use in the United States as a weight-control agent.

Dietary supplements can be marketed with no premarket safety evaluation by FDA. For dietary supplements that include an ingredient marketed in the United States before October 15, 1994—such as products containing sources of ephedrine alkaloids no FDA review is required. For dietary supplements that include an ingredient that was not marketed before October 15, 1994, manufacturers or distributors must submit a notice to FDA 75 days before marketing; however, the notice is not required to in-

*60 FR 38,643.

Ephedrine — Continued

clude objective evidence of safety, only an explanation of why there is a reasonable expectation that use of the supplement will be safe.

Because many of these products are marketed as "natural" or promoted as foods, consumers may assume incorrectly that the products are safe and without side effects. For example, the TDH investigation determined that, during medical evaluations, some patients did not report taking ephedrine-containing dietary supplements because they did not initially believe that a "natural" or "herbal" food supplement could be related to their illness. In addition, health-care providers or consumers may not have realized that ephedrine alkaloids and other stimulants were in the product because they were not included in the ingredient listing or because an unfamiliar name for the compound was used.

Because of misuse of and adverse reactions to products containing ephedrine, approximately 21 states have passed regulations stricter than federal regulations, including requiring that ephedrine drug and food products be made available by prescription only; moving ephedrine products to the schedules of controlled substances; and prohibiting weight loss, appetite control, or stimulant claims on the labels. Because of concerns about the safety of dietary supplements that contain sources of ephedrine alkaloids, a working group convened by FDA in October 1995 made several recommendations about potency limits and label warnings to promote safer use of these products. FDA has been evaluating these recommendations and, because of continuing concerns about the safety of these products, is convening a meeting of the Food Advisory Committee and the special working group on August 27–28, 1996, in Washington, D.C.

The findings in this report underscore the need for the general public and for health-care providers to be aware of potential health hazards associated with use of dietary supplements containing ephedrine and associated alkaloids. Health-care providers should question patients about their use of dietary supplements and herbal medications and report any adverse effects to dietary supplements, including those containing ephedrine and associated alkaloids, to FDA's MedWatch Program, telephone (800) 322-1088 ([800] FDA-1088). Consumers can report adverse events to the FDA Consumer Hotline, (800) 322-4010 ([800] FDA-4010).

References

- 1. Dollery C, ed. Ephedrine (hydrochloride). In: Therapeutic drugs. Vol 1. New York: Churchill Livingstone, 1991:E26–E29.
- 2. Pentel P. Toxicity of over-the-counter stimulants. JAMA 1984;252:1898–903.
- 3. Loizou LA, Hamilton JG, Tsementzis SA. Intracranial haemorrhage in association with pseudoephedrine overdose. J Neurol Neurosurg Psychiatry 1982;45:471–2.
- 4. Lake CR, Gallant S, Masson E, Miller P. Adverse drug effects attributed to phenylpropanolamine: a review of 142 case reports. Am J Med 1990;89:195–208.
- Breum L, Pedersen JK, Ahlstrom F, Frimodt-Moller J. Comparison of an ephedrine/caffeine combination and dexfenfluramine in the treatment of obesity: a double-blind, multi-centre trial in general practice. International Journal of Obesity and Related Metabolic Disorders 1994;18:99–103.
- 6. Astrup A, Lundsgaard C, Madsen J, Christensen NJ. Enhanced thermogenic responsiveness during chronic ephedrine treatment in man. Am J Clin Nutr 1985;42:83–94.
- Astrup A, Toubro S. Thermogenic, metabolic, and cardiovascular responses to ephedrine and caffeine in man. International Journal of Obesity and Related Metabolic Disorders 1993;17 (suppl):S41–S43.

Ephedrine — Continued

 Astrup A, Toubro S, Cannon S, Hein P, Breum L, Madsen J. Caffeine: a double-blind, placebocontrolled study of its thermogenic, metabolic, and cardiovascular effects in healthy volunteers. Am J Clin Nutr 1990;51:759–67.

Update: Diphtheria Epidemic — New Independent States of the Former Soviet Union, January 1995–March 1996

Epidemic diphtheria reemerged in the New Independent States (NIS) of the former Soviet Union, beginning in the Russian Federation in 1990 and affecting all 15 NIS by the end of 1994 (*1,2*). Approximately 90% of all diphtheria cases reported worldwide during 1990–1995 were reported from the NIS (World Health Organization [WHO], unpublished data, 1996). During 1993–1994, WHO, partner organizations, and national ministries of health developed a strategy to control the epidemic with a priority goal of achieving coverage of >90% among persons aged \geq 3 years with a single dose of diphtheria toxoid through mass vaccination campaigns and achieving coverage for routine childhood vaccination (i.e., four doses of diphtheria and tetanus toxoids and pertussis vaccine by age 2 years) of >95%. This report summarizes data provided to WHO about the incidence of diphtheria (Table 1) and efforts to implement control measures in the NIS during 1995 and January–March 1996.

Overall, from 1994 to 1995, reported diphtheria cases in the NIS increased 5.2%, from 47,628 to 50,412 cases (Figure 1) with approximately 1500 deaths in 1995. Since the epidemic began in 1990, approximately 125,000 cases and 4000 deaths have been reported in the NIS. The number of cases in 1995 reflects an 11% decrease in reported cases in the Russian Federation but a near doubling of cases from other NIS. However, following expanded control efforts, by the end of 1995 most countries began reporting decreases from the same periods in 1994. During January–March 1996, a total of 6179 diphtheria cases were reported in the NIS, a 59% decrease from the 14,931 cases reported during the same period in 1995.

Russian Federation. During 1990–1995, approximately 97,000 diphtheria cases and 2500 deaths were reported in the Russian Federation. Reported cases decreased from 39,582 (26.9 cases per 100,000 population) in 1994 to 35,652 (24.3) in 1995. Reported deaths decreased from 1104 (case-fatality rate: 2.8%) in 1994 to 740 (2.1%) in 1995. Vaccination coverage among persons aged \geq 18 years with at least one dose of diphtheria toxoid during the preceding 10 years had increased to 70%–80% by the end of 1995 from an estimated 20% in 1990. In 1995, reported coverage with a primary series of diphtheria toxoid among children aged 12–23 months was 92.7%.

Western NIS and Baltic countries. In Ukraine, reported cases nearly doubled, from 2990 (5.8 cases per 100,000) in 1994 to 5280 (10.3) in 1995. Mass vaccination campaigns targeting adults were conducted in seven of the 27 regions in April 1995; the remaining regions were targeted for mass vaccination in late 1995 and in 1996. Overall, vaccine coverage in adults was an estimated 60%. In Belarus, Latvia, and Lithuania, increases in diphtheria cases were reported in 1995 (Table 1); however, mass adult vaccination efforts in 1995 and 1996 achieved coverage of 60%–80%, and efforts to complete vaccination of all adults are continuing. In Moldova, reported cases increased from 376 (8.5 cases per 100,000 population) in 1994 to 418 (9.4) in 1995. A mass vaccination campaign was conducted in the summer and fall of 1995, increasing coverage among adults to >80%; a sustained decline in cases began in October 1995.

TABLE 1. Number and rate of diphtheria cases, 1994–1995, percentage change in number of cases from 1994 to 1995 and first
quarter of 1995 to first quarter of 1996, and level of coverage with diphtheria toxoid, 1995, by country — New Independent
States of the former Soviet Union

						% Change	in number of cases	% Coverage	, 1995	
		1994		1995		1994 to	1st quarter 1995 to	Primary series	1 dose,	
Country	Population*	Cases	Rate [†]	Cases	Rate	1995	1st quarter 1996	at age 1 year	adult	
Armenia	3.6	36	1.0	29	0.8	-19	-14	NA§	60¶	
Azerbaijan	7.5	841	11.3	883	11.7	5	-85	>90	95	
Belarus	10.1	230	2.3	322	3.2	40	-46	96	65	
Estonia	1.6	7	0.5	19	1.2	171	-29	>90	30**	
Georgia	5.4	294	5.4	419	7.7	43	-28	NA	65¶	
Kazakstan	17.0	489	2.9	1105	6.5	126	-31	93	50	
Kyrgyzstan	4.7	299	6.4	693	14.6	132	–15	93	70	
Latvia	2.6	250	9.7	369	14.4	48	-71	NA	30††	
Lithuania	3.6	38	1.0	43	1.2	13	-69	97	70	
Moldova	4.4	376	8.5	418	9.4	11	-85	NA	>80	
Russia	146.7	39,582	26.9	35,652	24.3	-10	-59	93	75	
Tajikistan	6.1	1912	31.8	4455	73.0	133	-42	94	>95	
Turkmenistan	4.1	60	1.5	87	2.1	45	90	92 ^{§§}	30	
Ukraine	51.3	2990	5.8	5280	10.3	77	-37	NA	60	
Uzbekistan	22.8	224	1.0	638	2.8	185	-40	NA	50	
Total	298.5	47,628	16.1	50,412	16.9	5	-59	_	_	

*In millions. [†]Per 100,000 population per year. [§]Not available.

[¶]Adult coverage for Armenia and Georgia is based on less complete data than for other countries. ** Approximately 75% of adults were vaccinated during 1985–1987. ^{††} The Latvian Ministry of Health considers coverage to be three doses for adults; an estimated 60%–70% have received at least one dose.

^{§§} Coverage figure is for four doses of diphtheria and tetanus toxoids and pertussis vaccine (primary series plus first revaccination) at age 2 years.

694

Diphtheria Epidemic — Continued





In Estonia, cases increased from seven to 19, almost exclusively among ethnic Russians in border areas. Approximately 30% of adults were vaccinated during the campaign in 1995.

Central Asia. All of the Central Asian republics reported increases in reported diphtheria cases in 1995. In Tajikistan, reported cases increased from 1912 (31.8 cases per 100,000 population) in 1994 to 4455 (73.0) in 1995, with cases reported from all regions of the country. A mass vaccination campaign was conducted during October 1995 and reached >90% of adults (aged ≤50 years). The other Central Asian republics (Kazakstan, Kyrgyzstan, Uzbekistan, and Turkmenistan) reported a total of 2523 cases in 1995 (Table 1), representing a 135% increase over the total reported cases in 1994. Mass vaccination campaigns targeting both adults and children were initiated in Kazakstan, Kyrgyzstan, and Uzbekistan in 1995 and are to be completed during 1996. In Turkmenistan, vaccination campaigns for children were conducted in 1995; adults are being targeted in 1996.

Caucasus. Georgia reported a 43% increase in diphtheria cases during 1995 (419) over 1994 (294), and Azerbaijan reported a 5% increase; Armenia continued to report few cases (Table 1). Following revaccination efforts among schoolchildren, resulting in coverage of 96%, Azerbaijan conducted mass vaccination of persons aged 0–55 years during November 1995, achieving coverage of 95%. Reported cases in Azerbaijan during January–March 1996 are 85% below the same period in 1995. In Georgia and Armenia, adult vaccination campaigns began in 1995; however, coverage is <60% in some regions.

Reported by: Regional Office for Europe, World Health Organization, Copenhagen, Denmark. International Federation of Red Cross and Red Crescent Societies, Geneva, Switzerland. Child

Diphtheria Epidemic — Continued

Vaccine Preventable Disease Br, Epidemiology and Surveillance Div, National Immunization Program, CDC.

Editorial Note: The findings in this report highlight the first evidence of progress toward controlling the diphtheria epidemic in the NIS. WHO continues to consider the epidemic an international public health emergency. At the time of the disintegration of the former Soviet Union in 1991, all NIS relied on supplies of vaccine and antitoxin from Russia, and most lacked the financial resources to procure them from the international market. Governmental and nongovernmental organizations, such as the United States Agency for International Development, European Community Humanitarian Office, International Federation of Red Cross and Red Crescent Societies, WHO, United Nations Children's Fund (UNICEF), the government of Japan, and others, have initiated an effort, monitored by the Interagency Immunization Coordination Committee, to mobilize the needed materials (i.e., vaccine, syringes, needles, antitoxin, and antibiotics) for all of the NIS except Russia, which continues to be self-sufficient for all these materials.

This epidemic has been characterized by a high proportion of cases among adults; children were the predominant age group affected in previous epidemics (3). Much of the variation in incidence rates in the NIS during 1994 and 1995 was attributed to differences in the timing of the onset of the epidemic between countries; however, the success in implementing control measures is beginning to affect incidence rates. In countries that have achieved high coverage among adults, diphtheria incidence has decreased substantially, regardless of the level of incidence before the vaccination campaigns. The impact of mass vaccination of the adult population on slowing the course of the epidemic underscores the need for rapidly completing the vaccination campaigns in the other countries.

All the NIS have attempted to increase childhood vaccination coverage, including decreasing perceived contraindications to childhood vaccination and increasing the routine use of full-strength vaccine preparations in the primary series; routine childhood coverage is now high in most countries. Most countries also have reinstituted a school-entry booster dose and many of the national mass campaigns have included preschool-aged and school-aged children and adolescents in the target population.

Other control measures are important in efforts to further reduce transmission of diphtheria. Most of the NIS have made efforts to improve early diagnosis and treatment of cases. Moldova, Lithuania, and Azerbaijan have adopted the WHO recommendation to use empiric antibiotic treatment for close contacts of persons with diphtheria. Providing additional doses of diphtheria toxoid to at least certain portions of the population (e.g., adults aged 30–50 years in whom risk for severe disease and death is highest) will be needed to fully protect all persons. Because some epidemiologic features of this epidemic differ from those of the prevaccine era and the epidemiologic situation may vary among the NIS, ongoing surveillance and additional field studies are needed to track the course of the epidemic in each country.

The reemergence of epidemic diphtheria throughout the NIS after >30 years of successful control and the reports of >20 imported cases from the NIS into Europe (1,2) and Mongolia (WHO, unpublished data, 1996) emphasize the need for achieving and maintaining high levels of diphtheria immunity among both adults and children in the United States and other countries. The Advisory Committee on Immunization Practices recommends that all children receive a routine series of five doses of diphtheria

Diphtheria Epidemic — Continued

toxoid-containing vaccine with doses at ages 2, 4, 6, and 12–20 months and 4–6 years; boosters of diphtheria and tetanus toxoids should then be administered beginning at age 11–12 years (provided at least 5 years have passed since the last dose of diphtheria toxoid-containing vaccine) and every 10 years thereafter (4–6). Travelers to areas with diphtheria activity should review their vaccination status and receive age-appropriate vaccinations as needed.

References

- 1. CDC. Diphtheria epidemic—New Independent States of the former Soviet Union, 1990–1994. MMWR 1995;44:177–81.
- 2. Hardy IR, Dittmann S, Sutter R. Current situation and control strategies for resurgence of diphtheria in Newly Independent States of the former Soviet Union. Lancet 1996;347:1739–44.
- 3. Stuart G. Note on diphtheria incidence in certain European countries. British Medical Journal 1945;2:613–5.
- 4. Advisory Committee and Immunization Practices. Diphtheria, tetanus, and pertussis: recommendations for vaccine use and other preventive measures—recommendations of the Immunization Practices Advisory Committee (ACIP). MMWR 1991;40(no. RR-10).
- 5. CDC. Recommended childhood immunization schedule—United States, January–June 1996. MMWR 1996;44:940–3.
- CDC. Food and Drug Administration approval of an acellular pertussis vaccine for the initial four doses of the diphtheria, tetanus, and pertussis vaccination series. MMWR 1996;45:676–7.

Progress Toward Poliomyelitis Eradication — Indonesia, 1995

In 1988, Indonesia (1991 population: 181 million) adopted the goal of eradicating poliomyelitis by the year 2000. Although routine coverage with three doses of oral poliovirus vaccine (OPV3) has been >90% among 1-year-old children since 1991, cases of polio continue to be reported (Figure 1). To interrupt poliovirus transmission, National Immunization Days (NIDs)* were conducted during September 13–17 and October 18–22, 1995, resulting in vaccination of >22 million children aged <5 years (representing approximately 100% of the target population). NIDs are planned for September 10–14 and October 15–19, 1996, and in 1997. This report presents the polio surveillance data for Indonesia for 1995, which indicate that substantial improvements are necessary to meet the objectives of the polio-eradication initiative. Activities are under way to intensify surveillance for acute flaccid paralysis (AFP) and to identify the remaining reservoirs of wild poliovirus transmission.

Indonesia, which instituted AFP surveillance in 1995, requires all cases of AFP to be reported immediately to health authorities. A case of suspected polio is defined as AFP, including Guillain-Barré syndrome, in a child aged <15 years for which no other cause can be immediately identified, or any patient in whom a clinician suspects polio regardless of age. Stool specimens collected from AFP case-patients were sent to one of three designated laboratories in Bandung, Jakarta, and Surabaya for virus isolation. Poliovirus isolates were sent to the national laboratory in Jakarta to distinguish wild and vaccine-type polioviruses. Genetic sequencing of wild polioviruses was conducted at CDC.

^{*}Mass campaigns over a short period (days to weeks) in which two doses of oral poliovirus vaccine are administered to all children in the target age group, regardless of prior vaccination history, with an interval of 4–6 weeks between doses.

Polio Eradication — Continued

FIGURE 1. Reported percentage coverage with three doses of oral poliovirus vaccine (OPV3) among 1-year-olds, by year, 1988–1994, and number of reported poliomyelitis cases, by year, 1988–1995 — Indonesia



Source: Ministry of Health, Indonesia.

The number of reported polio cases has decreased substantially since the start of the national Expanded Program on Immunization (Figure 1). In 1995, a total of 22 cases of AFP were reported, representing an AFP rate of 0.04 per 100,000 children aged <15 years; seven (33%) cases were investigated within 48 hours of report, 17 (78%) had stool specimens taken for virus isolation, and two (10%) cases had stool specimens taken within 2 weeks of paralysis onset. Of the 22 cases of suspected polio, 12 have been classified as confirmed polio cases (using the standard WHO case definition[†]) by virus isolation (five cases—two in case-patients and three in contacts of case-patients), with residual paralysis at 60 days' follow-up (four), or lost to follow-up (three). Four culture-confirmed cases of polio were associated with type 1 and one with type 3 wild polioviruses with onset of paralysis during March and August 1995. Four of these cases, in Java and Sumatra, occurred in unvaccinated children.

To define the molecular epidemiology of the type 1 wild polioviruses isolated from the four culture-confirmed cases in 1995, genetic sequencing was conducted on a 150nucleotide interval at the VP1/2A junction of the viral genome (1). Because the poliovirus genome mutates at a rate of approximately two nucleotide substitutions per week, it is possible to quantify the genetic relatedness of isolates from different geographic areas and determine chains of transmission. Data summarizing the sequence relatedness among wild poliovirus isolates recently identified as endemic in South and East Asia suggested that the type 1 wild polioviruses isolated from Indonesia in 1995 form a distinct cluster and have been circulating in the country for at least 10 years. The genetic sequences of the isolates from Java and Sumatra, although

[†]A confirmed case of polio is defined as AFP and at least one of the following: 1) laboratoryconfirmed wild poliovirus infection, 2) residual paralysis at 60 days, 3) death, or 4) no follow-up investigation at 60 days.

Polio Eradication — Continued

found at the same time on adjacent islands, differed enough to suggest that the two cases were not related epidemiologically.

Reported by: H Wibosono, MD, H Roespandi, MD, Surveillance Unit, Ministry of Health, Indonesia; W Gendro, Polio Laboratory, Indonesia National Institute of Health, Research, and Development, Jakarta; L Soemara, MD, PT Biofarma, S Rusliana, MD, Public Health Laboratory of Surabaya; S Prihatini, MD, Provincial Health Svcs, East Java. United Nations Children's Fund; World Health Organization, Jakarta, Indonesia. Respiratory and Enterovirus Br, National Center for Infectious Diseases; Polio Eradication Activity, National Immunization Program, CDC.

Editorial Note: Indonesia successfully conducted its first NIDs in 1995 and is improving surveillance for polio by requiring AFP reporting. The AFP reporting rate in Indonesia during 1995 (0.04 per 100,000) was substantially lower than the expected rate of one AFP case per 100,000 persons aged <15 years (the rate used to define a sensitive AFP surveillance system [653 cases would be expected each year in Indonesia if the rate reached one per 100,000]). To meet the objectives of the polio-eradication initiative, performance indicators other than the AFP rate (i.e., proportion of case investigations conducted within 48 hours of notification and proportion of cases for which two stool specimens were obtained within 2 weeks of paralysis onset) should be at least 80%, which indicates adequate surveillance. Major efforts are under way to strengthen the surveillance system as a guide to further polio-eradication activities in Indonesia; these efforts include establishing active AFP reporting (i.e., reviewing hospital discharge records for AFP in hospitalized patients) linked with timely AFP case investigations.

The low level of AFP reporting in Indonesia and the retrospective reviews of hospital records indicate that many persons with AFP or with physician-diagnosed polio are admitted to hospitals but are not reported to public health officials. Although a hospital early-warning system is in place to immediately report high-priority infectious diseases, active surveillance systems probably will be needed to increase sensitivity. These active systems will be valuable especially in densely populated urban areas.

For the global polio-eradication initiative, genetic analysis of wild poliovirus isolates has provided critically important information. The isolation of type 1 wild poliovirus in Java and Sumatra indicate that the virus is indigenous to Indonesia and has circulated for many years, and confirmed the independent evolution of at least two poliovirus reservoirs in the country. These findings suggest that wild poliovirus has remained endemic in Indonesia despite routine high coverage with OPV3 in recent years and underscore the need for supplementary vaccination strategies (i.e., NIDs) to interrupt poliovirus transmission.

Indonesia, the fourth most populous country in the world, is of critical importance to the global polio-eradication initiative. Improved virologic surveillance already has identified at least two indigenous poliovirus reservoirs. The government of Indonesia, in cooperation with the major partner agencies contributing to the polio-eradication initiative (including WHO, United Nations Children's Fund [UNICEF], and Rotary International), will need to establish a sensitive AFP surveillance system. Adequate surveillance is necessary to identify the remaining poliovirus reservoirs and target areas for supplemental vaccination activities (i.e., mopping-up vaccination[§]) and prepare Indonesia for eventual certification of polio-free status (2).

[§]House-to-house administration of two doses of oral poliovirus vaccine at an interval of 4–6 weeks to all children aged <3 years who reside in areas where risk for wild poliovirus transmission is highest.

Polio Eradication — Continued

References

- Kew OM, Mulders MN, Lipskaya GY, et al. Molecular epidemiology of polioviruses. Seminars in Virology 1995;6:401–14.
- World Health Organization. Report of the first meeting of the Global Commission for the Certification of the Eradication of Poliomyelitis. Geneva, Switzerland: World Health Organization, Global Programme for Vaccines and Immunization, Expanded Programme on Immunization, 1995; publication no. WHO/EPI/GEN/95.6.

Average Postpartum Length of Stay for Uncomplicated Deliveries — New Jersey, 1995

During 1970–1992, total lengths of hospital stay following delivery declined substantially for both mothers and newborn infants (*1*). In response to public perceptions that postpartum stays have become excessively short, 28 states have enacted legislation and the U.S. Senate is considering legislation* mandating that insurers provide health-care coverage for a minimum postpartum length of stay for both mother and infant. On June 28, 1995, New Jersey enacted legislation requiring insurance companies and health-maintenance organizations (HMOs) to pay for at least a 48-hour postpartum hospital stay for women and newborns following vaginal delivery and a 96-hour postpartum stay following cesarean delivery when requested by the mother or the attending physician.[†] To assess the effect of this law in New Jersey on the average length of stay for uncomplicated deliveries, electronic birth certificate (EBC) data for 1995 were analyzed from four New Jersey hospitals. This report summarizes the findings, which demonstrate that, following enactment of legislation, the average length of stay increased for uncomplicated deliveries at these four hospitals.

On January 17, 1995, New Jersey initiated pilot testing of an EBC system in four of the state's 73 facilities that perform deliveries (69 hospitals and four birthing centers). These four hospitals, representing approximately 8% of births in New Jersey, were selected because they were considered sufficiently diverse to test the system and to detect any problems before statewide implementation of the EBC system in all birthing facilities (V. Dato, Division of Family Health Services, New Jersey Department of Health and Senior Services, personal communication, September 1995). The selected hospitals (one regional perinatal center and three community hospitals) are nonprofit organizations and represent north, south, and central New Jersey and urban and sub-urban areas. Because the EBC data include the exact time of delivery and the day and hour of discharge for the infant, length of stay was calculated in hours and converted to the equivalent proportion of days.

Length of stay was defined as the time from birth to discharge of an infant from the hospital. Deliveries were classified as uncomplicated if they met the following criteria: 1) a live-born infant weighing \geq 2500 g (\geq 5 lbs 8 oz) was delivered vaginally or by cesarean, 2) the mother and infant were discharged on the same day, and 3) the infant was not admitted or transferred to a neonatal intensive-care unit. Because this study focused on length of stay for uncomplicated deliveries, deliveries with a high likeli-

^{*}The New Borns' and Mothers' Health Protection Act of 1996; S. Rep. No. 969, 104th Cong., 2d Sess. (1996).

[†]N.J. Stat. Ann. 17:48–61 as amended by Public Law 1995, c. 138.

Average Postpartum Length of Stay — Continued

hood of having complications (i.e., those greater than the 97.5th percentile for length of stay) were excluded from analysis.[§]

From January 17 through December 31, 1995, a total of 9007 live-born infants were delivered at the four selected hospitals in New Jersey. Of the 7209 uncomplicated deliveries, 5619 (78%) were vaginal, and 1590 (22%) were cesarean. The number of deliveries that occurred before and after the law were similar: for vaginal deliveries, 46% and 54% of deliveries, respectively, and for cesarean deliveries, 47% and 53%, respectively.

After enactment of the New Jersey legislation, the average postpartum length of stay increased for both uncomplicated vaginal and cesarean deliveries (Figure 1). The average length of stay for vaginal deliveries increased 29%, from 1.4 days before the law to 1.8 days after the law (Table 1). The average length of stay for cesarean deliveries increased 18%, from 2.8 days before the law to 3.3 days after the law.

Length of stay also was examined by type of insurance and hospital (Table 1). After enactment of the law, length of stay increased for all types of insurance; increases were greater for private and federal assistance insurance[¶] than for self-pay groups. However, self-pay deliveries represented <2% of all deliveries. Length of stay also increased for all four participating hospitals after enactment of the law; the increase was greatest for the hospital with the shortest average length of stay for cesarean deliveries before enactment of the law.

Reported by: V Dato, MD, L Ziskin, MD, M Fulcomer, PhD, RM Martin, MPH, K Knoblauch, New Jersey Dept of Health and Senior Svcs. Pregnancy and Infant Health Br, Div of Reproductive Health, National Center for Chronic Disease Prevention and Health Promotion, CDC.

Editorial Note: The findings in this report indicate that, after enactment of legislation in New Jersey mandating insurance coverage for a minimum length of hospital stay for mother and infant following obstetric delivery, the average postpartum length of stay for all uncomplicated deliveries increased by 10–12 hours in these four hospitals. Because the New Jersey EBC system allowed calculation of the length of stay in hours, the system provides a more precise measure of the length of stay than most previous studies, which calculated length of stay in days.

The findings in this report are subject to at least three limitations. First, EBC data are subject to the limitations of other vital records data, including limited information (particularly about content of prenatal care). Second, because the analysis included only the four hospitals using the EBC system for the 6 months before enactment of the New Jersey legislation, the results may not be generalizable to the entire state. Finally, some insurance carriers administer both traditional health-care plans, which are subject to state regulation, and self-insured health plans, which are not. Therefore, for deliveries covered by these carriers, the EBC system cannot differentiate those deliveries exempt from the state law.

Following consumer demand for less medical intervention in the birthing process (2) and institution of cost-containment measures by the insurance industry, the na-

[§]Deliveries for which the length of stay was greater than the 97.5th percentile for the four hospitals were identified using distributions specific for type of delivery and occurrence before or after enactment of the law. For vaginal deliveries, postpartum lengths of stay that were >2.5 days before the law and >2.8 days after the law were excluded from analysis and, for cesarean deliveries, postpartum lengths of stay >4.4 days before the law and >5.1 days after the law were excluded. The resulting distributions of length of stay for uncomplicated deliveries were approximately normal.

Includes Medicaid, Medicare, and other federal programs.

Average Postpartum Length of Stay - Continued

FIGURE 1. Average postpartum length of stay following uncomplicated deliveries*, by delivery method — four selected hospitals[†], New Jersey, January 17– December 31, 1995[§]



*Deliveries were classified as uncomplicated if they met the following criteria: 1) a live-born infant weighing ≥2500 g (≥5 lbs 8 oz) was delivered vaginally or by cesarean, 2) the mother and infant were discharged on the same day, and 3) the infant was not admitted or transferred to a neonatal intensive-care unit. Excluded from the analysis were deliveries with a length of stay greater than the 97.5th percentile for the four hospitals (i.e., those with a high likelihood of having complications). These were identified using distributions specific for type of delivery and occurrence before or after enactment of the law. For vaginal deliveries, postpartum lengths of stay that were >2.5 days before the law and >2.8 days after the law were excluded from analysis, and for cesarean deliveries, postpartum lengths of stay >4.4 days before the law and >5.1 days after the law were excluded. The resulting distributions of length of stay for uncomplicated deliveries were approximately normal.

[†]The selected facilities comprise one regional perinatal center and three community hospitals, are nonprofit organizations, and represent north, south, and central New Jersey and urban and suburban areas. These four of the state's 73 facilities that perform deliveries initiated pilot testing of an electronic birth certificate that includes the exact time of delivery and the day and hour of discharge for the infant, permitting calculation of length of stay in hours. [§]This study comprised 5619 vaginal deliveries and 1590 cesarean deliveries.

Source: 1995 New Jersey electronic birth certificate data.

		Vag	jinal		Cesarean						
Characteristic	Averag	e LOS	Absolute		Averag	le LOS	Absolute				
	Before law (n=2582)	After law (n=3037)	change (days)	Percentage change	Before law (n=754)	After law (n=836)	change (days)	Percentage change			
Mother's insurance [¶]											
Private**	1.4	1.8	0.4	29	2.8	3.3	0.5	18			
Federal assistance ^{††}	1.3	1.8	0.5	38	2.7	3.3	0.6	22			
Self-pay	1.3	1.7	0.4	31	2.8	3.0	0.2	7			
Hospital											
1	1.3	1.8	0.5	38	2.9	3.3	0.4	14			
2	1.3	1.9	0.6	46	2.3	3.1	0.8	35			
3	1.3	1.7	0.4	31	3.1	3.2	0.1	3			
4	1.5	1.9	0.4	27	3.0	3.5	0.5	17			
Total	1.4	1.8	0.4	29	2.8	3.3	0.5	18			

*In davs.

[†] Deliveries were classified as uncomplicated if they met the following criteria: 1) a live-born infant weighing \geq 2500 g (\geq 5 lbs 8 oz) was delivered vaginally or by cesarean, 2) the mother and infant were discharged on the same day, and 3) the infant was not admitted or transferred to a neonatal intensive-care unit. Excluded from the analysis were deliveries with a length of stay greater than the 97.5th percentile for the four hospitals (i.e., those with a high likelihood of having complications). These were identified using distributions specific for type of delivery and occurrence before or after enactment of the law. For vaginal deliveries, postpartum lengths of stay that were >2.5 days before the law and >2.8 days after the law were excluded from analysis, and for cesarean deliveries, postpartum lengths of stay >4.4 days before the law and >5.1 days after the law were excluded. The resulting distributions of length of stay for uncomplicated deliveries were approximately normal.

[§]The selected facilities comprise one regional perinatal center and three community hospitals, are nonprofit organizations, and represent north, south, and central New Jersey and urban and suburban areas. These four of the state's 73 facilities that perform deliveries initiated pilot testing of an electronic birth certificate that includes the exact time of delivery and the day and hour of discharge for the infant, permitting calculation of length of stay in hours.

[¶]Data were excluded for 1256 mothers (971 who had vaginal deliveries and 285 who had cesarean deliveries) whose insurance was categorized as "Other/Unknown insurance," and for 311 mothers (238 who had vaginal deliveries and 73 who had cesarean deliveries) for whom insurance information was missing.

** Includes commercial insurers, preferred-provider organizations, health-maintenance organizations (HMOs), and Medicaid HMOs.

^{††} Medicaid. Medicare, and other federal programs.

Average Postpartum Length of Stay — Continued

tional average length of hospital stay for obstetric deliveries declined by nearly 50% during 1970–1992. In 1992, the average total length of stay for all hospital deliveries in the United States was 2.6 days (2.1 days for vaginal deliveries and 4.0 days for cesarean deliveries) (1). In the western United States, lengths of stay of 12–24 hours after uncomplicated vaginal deliveries are common (3). However, consumer groups, legislators, and health-care providers are concerned that early hospital discharge may adversely affect 1) the mother's feeling of preparedness to care for the infant and herself, 2) the ability of hospital staff to effectively teach a mother to care for the infant and herself and assess her ability to do so, 3) initiation and continuation of breastfeeding, 4) early diagnosis and treatment of infant and maternal morbidity, and 5) the ability to obtain adequate specimens for newborn metabolic screenings (3).

Over the past several decades, early postpartum discharge has been variously defined (4). In 1992, the American Academy of Pediatrics (AAP) and the American College of Obstetricians and Gynecologists defined early discharge for uncomplicated deliveries as <48 hours for vaginal delivery and <96 hours for cesarean delivery (5). In addition, in 1995, the AAP published guidelines for newborn discharge (6) that include meeting certain medical criteria and receiving appropriate support and followup. However, previous studies have not adequately examined the effects of short postpartum stays on maternal and infant outcomes because of small sample sizes, poor study designs, and lack of controls. In addition, most of these studies were conducted under specific, controlled circumstances (e.g., intensive and repeated in-home follow-ups) that do not represent well-established standards of care for health-care providers (3). Further research is needed to determine an optimal length of stay and adequate follow-up care for women and infants.

Because of federal legislation enacted in 1974 (Employee Retirement Income Security Act of 1974 [ERISA]**), state legislation cannot mandate coverage requirements for employer self-insured plans. An estimated 40%–55% of the insured U.S. population is covered by ERISA plans (7). The proposed New Borns' and Mothers' Health Protection Act of 1996 would require a minimum coverage for postpartum stays of 48 hours for vaginal deliveries and 96 hours for cesarean deliveries for all insurance plans in all states.

References

- 1. CDC. Trends in length of stay for hospital deliveries—United States, 1970–1992. MMWR 1995;44:335–7.
- 2. Beck CT. Early postpartum discharge programs in the United States: a literature review and critique. Women Health 1991;17:125–38.
- 3. Braveman P, Egerter S, Pearl M, Marchi K, Miller C. Early discharge of newborns and mothers: a critical review of the literature. Pediatrics 1995;96:716–26.
- 4. Kessel W, Kiely M, Nora AH, Sumaya CV. Early discharge: in the end, it is judgement. Pediatrics 1995;96:739–42.
- American Academy of Pediatrics/American College of Obstetricians and Gynecologists. Guidelines for perinatal care. 3rd ed. Washington, DC: American College of Obstetricians and Gynecologists, 1992;105–8.
- Committee on Fetus and Newborn, American Academy of Pediatrics. Hospital stay for healthy term newborns. Pediatrics 1995;96:788–90.
- Kent C. Bill would put the brakes on "drive-through deliveries". American Medical News, October 2, 1995:1,19.

^{**} Public Law 93-406 (29 USC 1001–1461).



FIGURE I. Selected notifiable disease reports, comparison of 4-week totals ending August 10, 1996, with historical data — United States

*Ratio of current 4-week total to mean of 15 4-week totals (from previous, comparable, and subsequent 4-week periods for the past 5 years). The point where the hatched area begins is based on the mean and two standard deviations of these 4-week totals.

	Cum. 1996		Cum. 1996
Anthrax Brucellosis Cholera Congenital rubella syndrome Cryptosporidiosis* Diphtheria Encephalitis: California* eastern equine* St. Louis* western equine*	53 2 1 1,075 1 16 2 -	HIV infection, pediatric* [§] Plague Poliomyelitis, paralytic [¶] Psittacosis Rabies, human Rocky Mountain spotted fever (RMSF) Streptococcal toxic-shock syndrome* Syphilis, congenital** Tetanus Toxic-shock syndrome	170 - 22 357 10 - 17 89
Hansen Disease Hantavirus pulmonary syndrome* [†]	63 9	Trichinosis Typhoid fever	14 200

TABLE I. Summary — cases of selected notifiable diseases, United States, cumulative, week ending August 10, 1996 (32nd Week)

-: no reported cases

-: no reported cases *Not notifiable in all states. ¹ Updated weekly from reports to the Division of Viral and Rickettsial Diseases, National Center for Infectious Diseases (NCID). ⁵ Updated monthly to the Division of HIV/AIDS Prevention, National Center for HIV, STD, and TB Prevention (NCHSTP), last update July 30, 1996. [¶] Three suspected cases of polio with onset in 1996 have been reported to date. **Updated quarterly from reports to the Division of STD Prevention, NCHSTP. First quarter 1996 is not yet available.

				Esche	richia							
	AID	DS*	Chlamydia	COIL O	157:H7 PHLIS [§]	Gono	rrhea	Hep C/N	atitis A,NB	Legior	ellosis	
Reporting Area	Cum. 1996	Cum. 1995	Cum. 1996	Cum. 1996	Cum. 1996	Cum. 1996	Cum. 1995	Cum. 1996	Cum. 1995	Cum. 1996	Cum. 1995	
UNITED STATES	39,982	42,926	200,725	1,220	560	171,884	238,200	2,118	2,418	472	733	
NEW ENGLAND	1,589	2,182	10,662	177	33	4,480	4,582	72	81	25	15	
N.H.	29 50	70	397	10	12	29 80	52 71	6	12	-	4	
Vt.	14	21	-	12	11	37	34	27	7	3	-	
R.I.	113	996 144	4,068	/9	-	314	304	33 6	58 4	7	9 1	
Conn.	643	876	4,413	43	-	2,674	2,474	-	-	Ν	Ν	
MID. ATLANTIC	11,159	11,231	23,236	115	34	17,736	27,041	186	266	101	121	
N.Y. City	6,277	5,655	9,512	6	-	3,094 4,931	10,942	152	134	30	32	
N.J.	2,130	2,745	2,482	33	5	2,853	2,226	-	107	8	19	
Fa.	3 2 2 5	3 350	25,882	201	17	0,200 25,813	0,019 17 684	33 294	24 196	04 131	0/ 212	
Ohio	696	723	11,986	75	36	8,836	14,972	234	7	55	98	
Ind.	433	335	6,187	32	23	4,041	5,703	7	1	29	49	
Mich.	528	668	2,502 U	54	38	10,460 U	10,992	219	130	29	22	
Wis.	171	230	5,207	N	19	2,450	4,057	-	-	9	22	
W.N. CENTRAL	935 170	1,017 218	14,527	268 101	163 94	7,091	12,345	80 1	48	26	49	
lowa	63	54	2,588	66	50	663	798	40	8	6	16	
Mo. N Dak	469	472	7,640	38	- 7	4,927 1	7,049	21	15	6	13	
S. Dak.	8	9	689	9	-	95	120	-	1	2	-	
Nebr.	65 150	75	898	21	3	161	707	4	10	7	11	
S ATLANTIC	9 735	10 759	32 946	24 64	34	59 226	65 849	14	0 149	2 86	0 115	
Del.	193	219	1,148	-	1	875	1,303	1	-	8	2	
Md.	1,149	1,416 640	3,813 N	N	5	8,127 2 758	7,471	1	6	15	20	
Va.	647	880	6,532	Ν	16	5,800	6,437	8	9	13	11	
W. Va.	73 520	62	1	N 17	2	313	470	7	36	1	3	
S.C.	500	569	-	6	3	6,743	7,710	17	15	4	23	
Ga.	1,421	1,459 4 927	7,137	20 14	-	12,366	12,269	U 83	15 32	3	14 15	
ES CENTRAL	1,311	1,391	18 268	32	29	19 564	24 817	394	699	31	44	
Ky.	212	179	4,083	7	4	2,518	2,825	20	22	3	8	
Tenn. Ala	497 365	561 375	7,998 5,289	13	22	6,924 8 5 1 5	8,379 10 290	303 4	675 2	15	21 5	
Miss.	237	276	U	4	-	1,607	3,323	67	Ū	10	10	
W.S. CENTRAL	3,970	3,694	29,081	35	9	21,281	33,400	288	168	15	13	
Ark. La.	170 923	166 602	4.264	11 5	3	2,263	3,089	3 130	4 104	- 1	5 2	
Okla.	165	173	4,703	6	1	2,926	3,343	69	30	4	3	
	2,/12	2,753	20,114	13	2	11,375	19,503	206	30	10	3	
Mont.	22	1,330	9,073	12	40	4,480	5,578	12	289	24 1	4	
Idaho	25	31	944	20	5	67	86	88	34	-	2	
Colo.	335	454	350	34	21	1,077	33 1,797	35	42	3 7	33	
N. Mex.	114	111	U	5	-	540	636	42	35	1	4	
Utah	342 117	350 87	3,934 921	N 12	- 13	2,287	2,024	44 44	24 10	8	12	
Nev.	240	273	1,031	8	7	294	823	9	14	2	17	
PACIFIC	6,859	7,972	36,450	147	78 5	12,213	16,904	261	522	33	77	
Oreg.	311	275	5,855 U	47	29	331	468	4	33	-	-	
Calif.	5,964	6,910	25,872	66	36	10,105	14,059	99	346	28	57	
Hawaii	121	161	780	N	6	204	377	120	9	1	5	
Guam	4	-	168	Ν	-	31	77	1	4	2	1	
P.R. VI	1,352 16	1,692 25	N	12 N	U	157	363	71	141	-	-	
Amer. Samoa	-	-	-	Ň	Ŭ	-	15	-	-	-	-	
C.N.M.I.	1	-	N	N	U	11	32	-	5	-	-	

TABLE II. Cases of selected notifiable diseases, United States, weeks ending August 10, 1996, and August 12, 1995 (32nd Week)

N: Not notifiable U: Unavailable -: no reported cases C.N.M.I.: Commonwealth of Northern Mariana Islands

*Updated monthly to the Division of HIV/AIDS Prevention, National Center for HIV, STD, and TB Prevention, last update July 30, 1996. [†]National Electronic Telecommunications System for Surveillance. [§]Public Health Laboratory Information System.

	Lyı Dise	me ase	Mal	aria	Mening Dise	ococcal ase	Syp	hilis Secondary)	Tuberc	ulosis	Babies	Animal
Reporting Area	Cum.	Cum. 1995	Cum.	Cum. 1995	Cum.	Cum. 1995	Cum.	Cum. 1995	Cum. 1996	Cum. 1995	Cum.	Cum.
UNITED STATES	5,310	6,015	794	705	2,216	2,073	6,663	9,988	11,118	12,242	3,413	4,862
NEW ENGLAND Maine N.H. Vt. Mass. R.I.	1,817 13 21 9 103 204	1,221 10 18 7 70 189	33 6 1 2 11 5	29 3 1 9 3	96 12 3 3 36 10	96 6 16 6 33 4	109 1 50 1	230 2 1 39 2	242 4 8 1 113 24	295 11 9 2 162 28	425 56 41 106 65 30	988 21 109 123 314 195
Conn.	1,467	927	8	12	32	31	57	186	92	83	127	226
MID. ATLANTIC Upstate N.Y. N.Y. City N.J. Pa.	2,935 1,863 182 211 679	3,883 1,908 292 1,029 654	181 52 91 28 10	190 37 94 45 14	196 59 30 53 54	268 74 36 66 92	259 47 71 77 64	533 54 231 114 134	1,936 234 1,068 433 201	2,642 298 1,518 442 384	448 241 - 84 123	1,262 742 - 236 284
E.N. CENTRAL Ohio Ind. III. Mich. Wis.	38 25 12 1 U	253 20 11 13 5 204	85 9 35 22 10	97 6 12 54 13 12	300 118 46 77 31 28	297 86 41 80 54 36	851 310 139 284 U 118	1,714 546 184 678 177 129	1,200 176 112 689 161 62	1,149 161 109 610 217 52	40 6 1 7 15 11	50 5 9 6 22 8
W.N. CENTRAL Minn. Iowa Mo. N. Dak. S. Dak.	83 18 16 22	68 5 7 35 -	27 9 2 7 1	18 3 2 6 1 1	179 23 36 74 3 8	123 21 23 46 1 5	221 27 13 159 -	497 29 28 422	298 70 43 127 3 14	368 93 44 135 3 13	339 18 165 15 46 76	242 11 88 23 22 66
Nebr. Kans.	26	4 17	3 5	3	16	10	6 16	9	13	63	3 16	4 28
S. ATLANTIC Del. Md. D.C. Va. W. Va. N.C. S.C. Ga. Fla	272 36 138 1 25 9 43 3 1 16	413 30 278 2 30 18 35 9 8 3	176 3 7 24 3 15 9 16 60	135 1 37 11 30 1 11 - 14 30	484 2 48 35 11 58 44 111 167	341 5 29 45 8 57 44 66 83	2,282 23 368 98 279 1 643 254 381 235	2,522 8 269 73 380 8 709 376 470 229	2,064 20 186 82 178 40 301 221 395 641	2,169 37 242 65 146 51 255 204 405 764	1,686 43 399 8 349 67 438 57 191 134	1,320 70 265 10 253 77 306 95 175 69
E.S. CENTRAL Ky. Tenn. Ala. Miss.	42 8 16 5 13	35 8 17 2 8	18 3 8 3 4	12 1 4 5 2	125 20 16 50 39	136 35 48 29 24	1,489 83 525 371 510	2,002 113 507 395 987	812 155 237 269 151	850 186 280 247 137	133 32 45 54 2	174 15 65 90 4
W.S. CENTRAL Ark. La. Okla. Tex.	70 19 1 5 45	72 6 3 28 35	20 - 2 - 18	17 2 2 1 12	244 28 45 23 148	248 25 39 25 159	1,056 113 343 123 477	1,966 301 657 116 892	1,434 118 59 116 1,141	1,500 146 148 128 1,078	41 14 13 14 U	495 33 22 26 414
MOUNTAIN Mont. Idaho Wyo. Colo. N. Mex. Ariz. Utab	5 - 2 - - - 1	6 - 3 - 1 -	35 5 3 16 1 4	39 3 1 17 4 5	123 4 19 3 22 21 33 12	153 2 7 5 40 29 45 12	89 2 23 1 56 2	147 4 - 85 5 21 4	350 14 6 3 45 52 146 34	395 10 8 1 37 56 194 19	81 15 20 22 3 16 2	91 30 - 21 - 3 27 7
Nev.	-	2	2	3	9	13	3	28	50	70	3	3
PACIFIC Wash. Oreg. Calif. Alaska Hawaii	48 5 9 33 - 1	64 9 51 -	219 13 15 182 3 6	168 14 11 132 1 10	469 67 82 312 5 3	411 69 73 259 6 4	307 4 8 294 - 1	377 10 18 348 1 -	2,782 144 57 2,436 43 102	2,874 167 74 2,477 47 109	220 - 212 8 -	240 4 1 228 7 -
Guam P.R.	-	-	-	1 1	1 4	2 18	3 81	8 173	35 63	72 120	- 31	- 32
V.I. Amer. Samoa C.N.M.I.	-	-	-	2 - 1	-	-	- - 1	- - 1	-	3 23	- -	-

TABLE II. (Cont'd.) Cases of selected notifiable diseases, United States, weeks ending August 10, 1996, and August 12, 1995 (32nd Week)

N: Not notifiable U: Unavailable -: no reported cases

	H. influ	uenzae,		Hepatitis (vi	iral), by type			Measles	es (Rubeola)				
	inva	sive		A		В	Ind	igenous	lm	ported [†]			
Reporting Area	Cum. 1996*	Cum. 1995	Cum. 1996	Cum. 1995	Cum. 1996	Cum. 1995	1996	Cum. 1996	1996	Cum. 1996			
UNITED STATES	755	747	16,259	17,197	5,706	6,192	37	367	4	29			
NEW ENGLAND	20	30	199	163	110	151	-	8	1	4			
Maine N.H.	- 8	3 7	13 10	17 7	2 9	6 16	-	-	-	-			
Vt.	1	2	4	4	10	2	-	1	1	1			
Mass. R.I.	10 1	10 3	103	68 20	35	56 8	-	6	-	- 3			
Conn.	-	5	60	47	47	63	-	1	-	-			
MID. ATLANTIC	117 27	104	980 265	1,068	858	875	1	19	-	5			
N.Y. City	20	26	378	527	396	288	1	9	-	3			
N.J. Pa.	36 24	11 42	205 132	146 144	155 82	223 137	-	- 10	-	- 2			
E.N. CENTRAL	116	133	1,354	2,052	586	710	-	6	-	3			
Ohio	70	68 17	536	1,166	85 102	78	-	2	-	-			
III.	27	30	262	419	129	187	-	2	-	1			
Mich. Wis	7	16 2	264 100	235 134	232 38	256 50	-	1	-	2			
W.N. CENTRAL	33	55	1,314	1,192	260	418	-	17	1	2			
Minn.	20	28	77	118	35	33	-	14	1	2			
Mo.	5	17	621	853	127	304	-	2	-	-			
N. Dak. S. Dak.	- 1	- 1	28 37	19 33	-	4	-	-	-	-			
Nebr.	1	3	144	31	16	20	-	-	-	-			
	176	3 150	740	601	23	24	-	1	-	-			
Del.	2	- 150	9	8	900 6	6	-	1	-	-			
Md. D.C.	42 5	53	127 20	128 16	191 27	161 14	-	2	-	1			
Va.	6	19	99	118	90	65	-	-	-	2			
vv. va. N.C.	6 20	6 24	12 87	73	227	34 193	-	- 3	-	- 1			
S.C.	4	1 42	37	29 50	49	33	-	-	-	-			
Fla.	19	43	297	257	288	248	-	-	-	-			
E.S. CENTRAL	19	6	925	1,033	495	566	-	-	-	-			
Ky. Tenn.	4 8	-	628	33 847	285	441	-	-	-	-			
Ala. Miss	6 1	4	125 153	55 98	39 135	74	-	-	-	-			
W.S. CENTRAL	31	39	3.373	2.070	756	735	3	23	-	2			
Ark.		5	314	265	50	34	-		-	-			
Okla.	25	20	1,398	510	75 59	97	-	-	-	-			
Tex.	3	13	1,554	1,234	572	486	3	23	-	2			
MOUNTAIN Mont.	76	84	2,574 80	2,604 69	662 6	538 16	33	145	2	5			
Idaho	1	2	148	225	67	64	-	1	-	-			
Colo.	35 11	4 10	20	321	29 82	78	-	4	-	3			
N. Mex. Ariz	9	12 21	271 1 031	544 737	222 162	198 85	2	10 8	-	-			
Utah	6	9	585	494	65	46	31	117	2	2			
Nev.	5	26	155	131	29	34	-	5	-	-			
Wash.	2	140	321	6,324 494	60	1,363	-	45	-	-			
Oreg. Calif	22 140	20 114	562 3 828	1,610 4 084	39 963	83 1 161	-	4 30	-	- 2			
Alaska	1	-	30	27	9	9	-	63	-	-			
Hawaii	2	4	50	109	8	12	-	1	-	1			
P.R.	- 1	- 3	61	5 57	228	4 377	- -	- 7	-	-			
V.I. Amer Samoa	-	-	-	6 5	-	12	U	-	U	-			
C.N.M.I.	10	11	1	21	5	10	Ŭ	-	Ŭ	-			

TABLE III. Cases of selected notifiable diseases preventable by vaccination, United States, weeks ending August 10, 1996, and August 12, 1995 (32nd Week)

N: Not notifiable U: Unavailable -: no reported cases

*Of 175 cases among children aged <5 years, serotype was reported for 37 and of those, 10 were type b.

[†]For imported measles, cases include only those resulting from importation from other countries.

	Measles (Rul										
	Τα	otal		Mumps	5		Pertussi	s		Rubella	a
Reporting Area	Cum. 1996	Cum. 1995	1996	Cum. 1996	Cum. 1995	1996	Cum. 1996	Cum. 1995	1996	Cum. 1996	Cum. 1995
UNITED STATES	396	258	3	397	558	130	2,296	2,095	3	185	92
NEW ENGLAND	12	8	-	-	10	9	470	305	-	24	35
Maine	-	-	-	-	4	-	18	19	-	-	-
N.H. Vt	- 2	-	-	-	-	-	40 13	23	-	2	-
Mass.	9	2	-	-	2	9	394	211	-	20	7
R.I.	- 1	5	-	-	-	-	-	1	-	-	- 27
	24	11	-	-	ა იე	-	160	10	-	2	10
Upstate N.Y.	- 24	1	-	18	82 19	3	89	78	-	4	3
N.Y. City	12	5	-	13	9	-	21	27	-	1	7
N.J. Pa	- 12	5	-	2	13	- 7	5 54	52	-	2	2
E.N. CENTRAL	9	13	2	74	95	15	225	254	-	3	3
Ohio	2	1	2	32	29	9	110	79	-	-	-
Ind.	- 3	- 1	-	5 18	7 28	- 5	19	18 45	-	-	-
Mich.	3	5	-	18	31	1	22	35	-	2	3
Wis.	1	6	-	1	-	-	5	77	-	-	-
W.N. CENTRAL	19	2	-	9	32	34	127	124	-	1	-
lowa	10	-	-	3 1	2	30	89 4	42	-	- 1	-
Mo.	2	1	-	2	18	1	20	36	-	-	-
N. Dak. S. Dak	-	-	-	2	-	- 1	1 3	6	-	-	-
Nebr.	-	-	-	-	4	2	6	7	-	-	-
Kans.	1	1	-	1	-	-	4	19	-	-	-
S. ATLANTIC	11	11	1	65	85	9	290	173	-	89	8
Md.	3	- 1	-	19	27	6	105	21	-	-	1
D.C.	-	-	-	-	-	-	-	4	-	1	-
va. W. Va.	2 -	-	-	-	16	-	26	-	-	2 -	-
N.C.	4	-	-	14	16	-	49	81	-	75	1
S.C. Ga.	- 1	2	-	5	6	-	13	10	-	-	-
Fla.	-	8	-	15	13	1	62	19	-	10	6
E.S. CENTRAL	-	-	-	18	7	2	62	96	1	3	1
Ky. Tenn	-	-	-	- 1	-	-	26 17	11 55	- 1	- 1	- 1
Ala.	-	-	-	3	4	2	12	30	-	2	-
Miss.	-	-	-	14	3	-	7	-	N	N	N
W.S. CENTRAL	25	20	-	16	38	2	59 4	163 27	-	2	7
La.	-	18	-	11	8	-	6	11	-	1	-
Okla.	- 25	-	-	-	- 25	-	8	17	-	-	- 7
	150	68	_	22	25	13	2/2	/01		6	1
Mont.	-	-	-	-	1	1	12	3	-	-	-
Idaho	1	-	-	-	2	11	85	85	-	2	-
Colo.	7	26	-	2	-	1	64	64	-	2	-
N. Mex.	10	31	N	N	N	-	34	63	-	-	-
Utah	119	-	-	2	11	-	11	140	-	-	3
Nev.	5	1	-	17	9	-	22	22	-	1	-
PACIFIC	146	125	-	136	184	36	652	411	2	50	22
vvasn. Oreg.	45 4	19	-	18	10	-	235	96 27	-	1	-
Calif.	32	103	-	99	158	29	374	250	2	45	18
Alaska Hawaii	63 2	- 2	-	2 17	12 4	-	2 12	- 38	-	- 3	- 4
Guam	-	-	П	5	2	П	1	2	Ш	-	
P.R.	7	3		1	2		1	1		-	-
V.I. Amer Samoa	-	-	U	-	3	U	-	-	U	-	-
C.N.M.I.	-	-	Ŭ	-	-	Ŭ	-	-	Ŭ	-	-

TABLE III. (Cont'd.) Cases of selected notifiable diseases preventable by vaccination,United States, weeks ending August 10, 1996, and August 12, 1995 (32nd Week)

N: Not notifiable U: Unavailable -: no reported cases

	All Causes, By Age (Years)						P&I [†]			All Cau	ises, B	y Age (Y	'ears)		P&I [†]
Reporting Area	All Ages	>65	45-64	25-44	1-24	<1	Total	Reporting Area	All Ages	>65	45-64	25-44	1-24	<1	Total
NEW ENGLAND Boston, Mass. Bridgeport, Conn. Cambridge, Mass. Hartford, Conn. Lowell, Mass. Lynn, Mass. New Bedford, Mass. New Haven, Conn. Providence, R.I. Somerville, Mass. Springfield, Mass. Waterbury, Conn.	533 145 38 9 18 48 30 13 27 49 41 3 37 28 28	386 94 9 12 32 25 9 24 30 32 35 23	82 28 6 5 7 4 2 1 7 6 2 2 3 0	43 12 1 8 2 2 8 1 1 2	13 6 2 - 1 2 2 2	95	37 92 213 231 140	S. ATLANTIC Atlanta, Ga. Baltimore, Md. Charlotte, N.C. Jacksonville, Fla. Miami, Fla. Norfolk, Va. Richmond, Va. Savannah, Ga. St. Petersburg, Fla. Tampa, Fla. Washington, Dcl. Wilmington, Del. F.S. CENTRAI	1,388 176 272 67 116 116 52 67 38 36 159 273 16 774	821 92 158 47 72 70 30 42 27 27 104 146 6 497	307 49 55 9 22 25 12 16 6 9 33 61 10	176 21 42 7 19 15 3 6 4 - 17 42 - 60	46 7 8 2 3 4 1 1 - 2 15 - 21	36 7 9 2 - 3 3 1 - 2 9 -	60 1 26 3 2 2 3 3 3 3 - 11 6 - 59
MiD. ATLANTIC Albany, N.Y. Allentown, Pa. Buffalo, N.Y. Camden, N.J. Elizabeth, N.J. Erie, Pa.§	47 2,313 45 15 85 30 10 33	32 1,516 31 12 71 13 5 29	9 455 2 10 11 3 4	5 259 6 1 2 2 1	46 3 - 1 -	37 - 1 4 1	9 101 3 1 - 3	Birmingham, Ala. Chattanooga, Tenn. Knoxville, Tenn. Lexington, Ky. Memphis, Tenn. Mobile, Ala. Montgomery, Ala. Nashville, Tenn.	103 92 94 69 182 77 28 129	59 60 73 39 123 46 16 81	26 21 12 21 42 15 8 26	7 6 4 5 12 8 2 16	4 1 5 - 5 4 1	25 4 - 4 - 4 1 5	4 8 11 4 22 4 - 6
Jersey City, N.J. New York City, N.Y. Newark, N.J. Paterson, N.J. Philadelphia, Pa. Pittsburgh, Pa.§ Reading, Pa. Rochester, N.Y. Schenectady, N.Y. Scranton, Pa.§ Syracuse, N.Y. Trenton, N.J. Utica, N.Y. Yonkers, N.Y.	35 1,152 65 34 400 46 15 139 10 25 109 20 19 26	22 743 29 11 250 32 12 99 6 21 84 11 16 19	7 230 21 9 87 5 - 28 3 3 15 4 2 6	4 144 10 13 44 5 2 9 1 1 8 4 1 1	2 18 3 - 1 2 - 1 - 1 -	17 2 1 4 4 - 1 1 1 1	3 24 4 13 5 3 19 1 10 1 5	W.S. CENTRAL Austin, Tex. Baton Rouge, La. Corpus Christi, Tex. Dallas, Tex. El Paso, Tex. Ft. Worth, Tex. Houston, Tex. Little Rock, Ark. New Orleans, La. San Antonio, Tex. Shreveport, La. Tulsa, Okla.	1,458 53 60 59 178 77 121 322 66 138 180 94 110	948 38 37 105 55 212 47 93 118 62 68	280 7 10 16 30 14 23 67 12 23 34 19 25	146 5 29 4 15 27 4 17 15 6 9	44 2 1 4 3 5 9 2 3 8 3 2	40 1 10 1 3 7 1 2 5 4 6	75 5 3 4 5 29 2 2 8 12 4
E.N. CENTRAL Akron, Ohio Canton, Ohio Chicago, III. Cincinnati, Ohio Cleveland, Ohio Columbus, Ohio Dayton, Ohio Detroit, Mich. Evansville, Ind. Fort Wayne, Ind. Gary, Ind. Grand Rapids, Mich Indianapolis, Ind. Madison, Wis. Peoria, III. Rockford, III. South Bend, Ind. Toledo, Ohio Youngstown, Ohio	2,066 38 33 513 147 154 212 108 181 39 65 49 33 45 52 95 49	1,323 26 28 275 103 97 136 68 97 30 46 U 45 96 U 64 24 24 24 24 24 24 24 24 24 24 24 24 24	$\begin{array}{c} 397\\ 10\\ 3\\ 108\\ 256\\ 42\\ 31\\ 35\\ 5\\ 12\\ U\\ 10\\ 21\\ 5\\ 12\\ 4\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 5\\ 12\\ 12\\ 5\\ 12\\ 12\\ 5\\ 12\\ 12\\ 5\\ 12\\ 12\\ 12\\ 5\\ 12\\ 12\\ 12\\ 12\\ 12\\ 12\\ 12\\ 12\\ 12\\ 12$	191 1 694 14 12 6 35 2 3 U 5 9 U 8 4 2 2 2 2	87 - 30 6 13 2 6 13 2 11 1 4 9 U 1 - 3 1 -	65 1 28 3 1 9 1 3 1 - U 2 2 U 5 - - 8 - 8	106 2 34 10 3 13 8 - 1 5 U 5 3 U 4 3 4 4 6 1	MOUNTAIN Albuquerque, N.M. Colo. Springs, Colo Denver, Colo. Las Vegas, Nev. Ogden, Utah Phoenix, Ariz. Pueblo, Colo. Salt Lake City, Utah Tucson, Ariz. PACIFIC Berkeley, Calif. Fresno, Calif. Glendale, Calif. Honolulu, Hawaii Long Beach, Calif. Dos Angeles, Calif. Pasadena, Calif. Portland, Oreg. Sacramento, Calif. San Diego, Calif.	873 89 53 106 155 29 20 95 119 1,703 16 82 17 74 64 340 32 153 153 153	597 69 38 80 104 22 119 56 90 1,137 90 1,137 90 13 53 43 232 19 90 106 857	$153 \\ 12 \\ 9 \\ 10 \\ 252 \\ -0 \\ 22 \\ 320 \\ 4 \\ 14 \\ 24 \\ 13 \\ 72 \\ 6 \\ 28 \\ 21 \\ 21 \\ 28 \\ 21 \\ 20 \\ 14 \\ 28 \\ 21 \\ 20 \\ 21 \\ 20 \\ 21 \\ 20 \\ 21 \\ 20 \\ 21 \\ 20 \\ 21 \\ 20 \\ 21 \\ 21$	74 7 12 32 1 9 4 156 1 9 2 5 5 5 25 3 9 14 14	32 1 2 2 9 2 8 - 5 3 4 1 6 - 2 3 7 - 6 3 7	15 23 5 5 43 13 - 44 22 92	54 4 10 5 1 4 6 10 155 9 1 15 17 4 4 22 17
W.N. CENTRAL Des Moines, Iowa Duluth, Minn. Kansas City, Kans. Kansas City, Mo. Lincoln, Nebr. Minneapolis, Minn. Omaha, Nebr. St. Louis, Mo. St. Paul, Minn. Wichita, Kans.	710 U 35 49 108 26 131 73 117 58 113	486 U 26 37 60 19 95 58 77 46 68	119 U 6 23 4 19 9 23 8 22	43 U 3 2 10 1 8 2 6 3 8	26 U 3 1 4 2 4 1 10	25 U 2 3 1 5 2 7 5	24 U 2 3 8 - 6 2 - 2 1	San Francisco, Calif. San Jose, Calif. Santa Cruz, Calif. Seattle, Wash. Spokane, Wash. Tacoma, Wash. TOTAL	1. 144 167 21 154 67 113 11,818 ¹	87 110 14 97 45 85 7,711	30 37 4 27 15 17 2,284	21 12 3 24 3 6 1,148	1 3 1 3 359	3 7 3 2 293	15 2 3 5 9 671

TABLE IV. Deaths in 121 U.S. cities,* week ending August 10, 1996 (32nd Week)

U: Unavailable -: no reported cases *Mortality data in this table are voluntarily reported from 121 cities in the United States, most of which have populations of 100,000 or more. A death is reported by the place of its occurrence and by the week that the death certificate was filed. Fetal deaths are not included. *Pneumonia and influenza. *Because of changes in reporting methods in these 3 Pennsylvania cities, these numbers are partial counts for the current week. Complete counts will be available in 4 to 6 weeks. *Total includes unknown ages.

Contributors to the Production of the MMWR (Weekly)

Weekly Notifiable Disease Morbidity Data and 121 Cities Mortality Data

Denise Koo, M.D., M.P.H. Deborah A. Adams Timothy M. Copeland Patsy A. Hall Carol M. Knowles Sarah H. Landis Myra A. Montalbano

Desktop Publishing and Graphics Support

Jolene W. Altman Morie M. Higgins Peter M. Jenkins

The Morbidity and Mortality Weekly Report (MMWR) Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy on Friday of each week, send an e-mail message to *lists@list.cdc.gov*. The body content should read *subscribe mmwr-toc*. Electronic copy also is available from CDC's World-Wide Web server at *http://www.cdc.gov/* or from CDC's file transfer protocol server at *ftp.cdc.gov*. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone (202) 512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to: Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone (404) 332-4555.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

Director, Centers for I and Prevention David Satcher, M. Deputy Director, Cent and Prevention Claire V. Broome, Director, Epidemiolog Stephen B. Thacke Editor, MMWR Series	Disease Control D., Ph.D. ers for Disease Control M.D. y Program Office er, M.D., M.Sc.	Acting Editor, <i>MMWR</i> (weekly) Andrew R. Pelletier, M.D. Managing Editor, <i>MMWR</i> (weekly) Karen L. Foster, M.A. Writers-Editors, <i>MMWR</i> (weekly) David C. Johnson Darlene D. Rumph Person Caran R. Wilbanks Editorial Assistant, <i>MMWR</i> (weekly)	
Richard A. Goodm	ian, M.D., M.P.H.	Teresa F. Rutledge	
☆U.S	. Government Printing Offic	e: 1996-733-175/47020 Region IV	