



- **433** World No-Tobacco Day
- 433 Cigar Smoking Among Teenagers United States, Massachusetts, and New York, 1996
- 440 Illegal Sales of Cigarettes to Minors — Mexico City, Mexico, 1997
- **444** Smoking-Attributable Mortality and Years of Potential Life Lost — United States, 1984
- United States, 1984 451 Outbreaks of Cyclosporiasis — United States, 1997
- 452 Notice to Readers

World No-Tobacco Day — May 31, 1997

MORBIDITY AND MORTALITY WEEKLY REPORT

World No-Tobacco Day is an annual international event that encourages governments, communities, and other groups to become more aware of the hazards of tobacco use and requests all persons who use tobacco to quit for at least 24 hours. This year's event will be held May 31; the theme is "United for a Tobacco-Free World" (1).

Tobacco use is expected to be the greatest risk factor for death and disability in the world by 2020 (2). In 1990, approximately 3 million deaths were attributed to tobacco use; by 2025, the annual number of tobacco-related deaths is projected to reach 10 million, with 70% of deaths occurring in developing countries (1). Efforts to reduce tobacco use require the participation of all sectors of society and must be comprehensive in scope. This year's event will highlight the complementary roles of policies and programs at the local, national, and international levels in achieving a tobacco-free world.

The World Health Organization (WHO), which is sponsoring this year's event, will provide press releases, fact sheets, a poster, and an advisory kit on comprehensive measures to reduce tobacco use. Additional information is available from WHO on the Internet (http://www.who.ch/programmes/psa/toh.htm), the WHO Regional Office for the Americas (telephone [202] 974-3000), and from CDC's Office on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion (http://www.cdc.gov/tobacco) (telephone [770] 488-5705).

References

- 1. World Health Organization. World No-Tobacco Day, 31 May 1997 [Advisory kit]. Geneva: World Health Organization, 1996.
- 2. World Health Organization. Investing in health research and development. Geneva: World Health Organization, 1996.

Cigar Smoking Among Teenagers — United States, Massachusetts, and New York, 1996

Cigar smoking can cause cancers of the oral cavity, larynx, esophagus, and lung (1) and chronic obstructive pulmonary disease (2). In addition, cigars contain substantial levels of nicotine, an addictive drug (3). Despite these health risks, total cigar consumption in the United States was approximately 4.5 billion cigars in 1996, and con-

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES / Public Health Service

Cigar Smoking Among Teenagers — Continued

sumption of larger cigars increased by 44.5% from 1993 through 1996 (from 2,138 million cigars to 3,090 million cigars, respectively) (4). This report presents estimates of the prevalence of cigar smoking among youth based on analyses of data from the Robert Wood Johnson Foundation's (RWJF) 1996 National Study of Tobacco Price Sensitivity, Behavior, and Attitudes Among Teenagers and Young Adults; a 1996 survey by the Massachusetts Department of Public Health (MDPH) of high school and junior high school students; and the Roswell Park Cancer Institute's 1996 Survey of Alcohol, Tobacco, and Drug Use in two New York counties (*5*). The analyses indicate that, during the year before being surveyed, 26.7% of U.S. and 28.1% of Massachusetts high school students in two New York counties reported having smoked cigars during the previous 30 days.

National Survey

The RWJF survey employed a three-stage cluster sample design to produce a nationally representative sample of students in grades 9-12. Within the selected sample of 200 counties (primary sampling units), schools were randomly selected, with the probability of selection proportional to enrollment size. Four alternate high schools were simultaneously selected, matching the original school in size, type, location, and the race/ethnicity and socioeconomic status of the students. An alternate was substituted when the first school chosen for the study could not participate. A total of 202 schools (representing 146 [73%] of the 200 primary sampling units) participated in the study. Within each school, one class per grade was chosen randomly. All students in the selected classes were eligible to participate; 80% of the students enrolled in the sample of selected classes participated. A total of 16,556 students aged 14–19 years completed the survey; however, 139 were excluded from these analyses because of missing information on sex. Participants were asked, "How many cigars, if any, have you smoked in the past year?" Annual cigar smokers were defined as any student who reported smoking a cigar during the previous year; frequent cigar smokers were defined as any student who reported smoking \geq 50 cigars during the previous year. Data were weighted by age, race/ethnicity, sex, and region* to provide national estimates. Confidence intervals (CIs) were calculated using SUDAAN.

In 1996, an estimated 6.0 million (26.7% [95% Cl= \pm 1.7%]) 14–19-year-olds reported having smoked a cigar during the previous year (4.3 million [37.0% (95% Cl= \pm 2.4%)] males and 1.7 million [16.0% (95% Cl=1.3%)] females) (Table 1). Cigarette smokers were more than three times as likely as noncigarette smokers to report having smoked a cigar (54.1% [95% Cl= \pm 2.4%], compared with 14.2% [95% Cl= \pm 1.2%], respectively). Among the 68.8% of students who did not smoke cigarettes, males were more likely than females to have reported smoking a cigar during the previous year (20.4% [95% Cl= \pm 1.8%] versus 7.8% [95% Cl= \pm 1.1%], respectively). Users of smokeless tobacco were more than three times as likely as nonusers to report having smoked cigars

^{*}The four regions were Northeast (Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont), *Midwest* (Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin), *South* (Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia), and *West* (Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming).

TABLE 1. Percentage of students aged 14–19 years who reported having smoked at least one cigar during the previous yea
by selected characteristics — United States, National Study of Tobacco Price Sensitivity, Behavior, and Attitudes Amon
Teenagers and Young Adults, 1996

			Annual	cigar use*				Fr	equen	t cigar use†		
	F	emale		Male	1	Total	F	emale		Male	1	fotal
Characteristic	%	(95% Cl [§])	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)	%	(95% CI)
Race/Ethnicity												u U
White, non-Hispanic	16.0	(± 1.8%)	41.6	(±2.7%)	28.9	(± 2.1%)	1.2	(±0.4%)	3.4	(±0.8%)	2.3	(±0.5%)
Black, non-Hispanic	13.4	(± 2.9%)	25.2	(±3.7%)	19.3	(± 2.9%)	1.6	(±0.6%)	5.6	(±2.0%)	3.6	(±1.1%)
Hispanic	20.0	(± 3.0%)	32.3	(±3.0%)	26.2	(± 2.1%)	1.8	(±0.8%)	3.2	(±0.9%)	2.5	(±0.6%) थ्रे
Other [¶]	14.5	(± 2.7%)	28.5	(±4.3%)	22.2	(± 2.9%)	0.5	(±0.7%)	5.8	(±2.4%)	3.4	(±1.3%) है
Age group (yrs)												
14–16	16.8	(± 1.8%)	32.1	(± 2 .4%)	24.4	(±1.7%)	1.3	(±0.4%)	2.9	(±0.7%)	2.1	(±0.4%) g
17–18	14.9	(± 1.8%)	43.9	(±3.2%)	29.8	(± 2.4%)	1.1	(±0.5%)	5.2	(±1.1%)	3.2	(±0.7%)
19	14.9	(± 6.7%)	35.5	(±7.8%)	27.5	(±5.3%)	3.1	(±3.0%)	4.9	(±2.6%)	4.2	(±2.1%)
Region**												
Northeast	12.4	(± 2.1%)	33.7	(±5.2%)	23.2	(± 3.4%)	0.8	(±0.5%)	3.0	(±1.0%)	1.9	(±0.5%)
Midwest	16.9	(± 2.6%)	42.3	(±4.2%)	29.8	(± 3.4%)	1.3	(±0.7%)	4.9	(±1.5%)	3.2	(±0.9%)
South	17.2	(± 2.2%)	37.1	(±3.7%)	27.3	(± 2.4%)	1.5	(±0.5%)	4.3	(±1.3%)	3.0	(±0.9%)
West	16.4	(± 3.4%)	34.5	(±5.4%)	25.6	(± 4.2%)	1.1	(±0.5%)	3.0	(±1.0%)	2.1	(±0.5%)
Education of parents ^{††}												
Completed college	16.0	(± 2.0%)	38.7	(±3.1%)	27.9	(± 2.2%)	1.0	(±0.4%)	3.5	(±0.9%)	2.3	(±0.5%)
Did not complete college	16.0	(± 1.5%)	37.0	(±3.3%)	26.1	(± 2.1%)	1.3	(±0.4%)	4.0	(±0.8%)	2.6	(±0.5%)
School performance												
Better or much better												
than average	12.5	(± 1.6%)	31.2	(±2.9%)	21.5	(± 2.0%)	0.7	(±0.3%)	3.2	(±0.8%)	1.9	(± 0.4%)
Average	19.2	(± 1.8%)	40.9	(±3.0%)	30.1	(±2.1%)	1.6	(±0.5%)	4.3	(±1.0%)	3.0	(±0.6%)
Below average	28.6	(± 6.0%)	54.7	(±5.4%)	45.1	(± 4.2%)	3.5	(±1.9%)	5.4	(±2.1%)	4.7	(±1.6%)
Household smoker												
Present	19.1	(± 1.7%)	41.2	(±3.1%)	30.1	(±2.1%)	1.8	(±0.5%)	4.9	(±1.1%)	3.4	(±0.6%)
Not present	13.0	(± 1.7%)	33.0	(±2.8%)	23.2	(±1.9%)	0.7	(±0.4%)	3.0	(±0.6%)	1.8	(± 0.4%)
Cigarette use												
Smoker ^{§§}	34.0	(± 2.6%)	73.9	(±2.6%)	54.1	(± 2.4%)	3.0	(±0.8%)	9.4	(±1.7%)	6.2	(±0.9%)
Nonsmoker	7.8	(± 1.1%)	20.4	(±1.8%)	14.2	(±1.2%)	0.3	(±0.2%)	1.5	(±0.4%)	0.9	(±0.2%)

Vol. 46 / No. 20 Cigar Smoking / r` g l

MMWR

435

Smokeless tobacco use User¶¶ Nonuser	50.1 15.5	(±12.0%) (± 1.3%)	75.9 30.6	(±3.6%) (±1.9%)	73.4 22.6	(±3.4%) (±1.4%)	3.3 1.2	(±3.5%) (±0.3%)	8.4 3.2	(±2.4%) (±0.6%)	7.9 2.1	(±2.2%) (±0.3%)	Cigar Sn
Total	16.0	(± 1.3%)	37.0	(± 2.4%)	26.7	(±1.7%)	1.2	(±0.3%)	3.9	(±0.6%)	2.6	(± 0.4%)	nokii

*Smoked one or more cigars during the previous year.

[†]Smoked \geq 50 cigars during the previous year.

[§]Confidence interval.

[¶]Numbers for other races were too small for meaningful analysis.

** Northeast=Connecticut, Maine, Massachusetts, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, and Vermont; Midwest=Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin; South=Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, South Carolina, Tennessee, Texas, Virginia, and West Virginia; and West=Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming.

^{††} Highest level of education of either parent.

§§ Smoked during the previous 30 days.

"Used chewing tobacco or snuff during the previous 30 days.

Cigar Smoking Among Teenagers — Continued

(73.4% [95% Cl= \pm 3.4%], compared with 22.6% [95% Cl= \pm 1.4%], respectively). Cigar smoking did not vary substantially by region or race/ethnicity, although prevalence was greatest among white, non-Hispanic males (41.6% [95% Cl= \pm 2.7%]).

Massachusetts Survey

The MDPH survey sample comprised two subsamples of students in grades 6–12: a statewide random sample, proportionately stratified by area and grade, and a separate random sample of five urban areas in the state, stratified by percentage of non-white students in each grade. These five urban areas were selected to oversample communities with racial/ethnic minorities to ensure adequate representation for analysis. Of the 191 schools meeting eligibility criteria, 171 (90%) participated in this survey. Of the 8236 students eligible to participate in the survey, 6844 (83.1%) participated. Data were collected during November 1996–January 1997. School and class selection was random, participation was voluntary, and all responses were anonymous. The questionnaires were self-administered. All students were asked "How often have you smoked cigars in your lifetime?"; "How often have you smoked cigars during the last 12 months?"; and "How often have you smoked cigars during the last 30 days?" The response categories were never, one to two times, three to five times, six to nine times, 10–19 times, 20–39 times, and ≥40 times.

Among the 1020 students in grade 6, 9.9% (95% Cl= \pm 1.8%) reported having ever smoked a cigar, 5.0% (95% Cl= \pm 0.8%) smoked a cigar during the previous year, and 2.0% (95% Cl= \pm 0.9%) smoked a cigar during the previous month. Among 1942 students in grades 7 and 8, 22.3% (95% Cl= \pm 1.8%) reported having ever smoked a cigar, 14.1% (95% Cl= \pm 1.5%) smoked a cigar during the previous year, and 7.6% (95% Cl= \pm 1.2%) smoked a cigar during the previous month. Among the 3873 high school students in grades 9–12, 38.9% (95% Cl= \pm 1.5%) reported having ever smoked a cigar, 28.1% (95% Cl= \pm 1.4%) smoked a cigar during the previous year, and 14.5% (95% Cl= \pm 1.1%) smoked a cigar during the previous month.

High school students who had used other tobacco products during the previous month were also more likely to have smoked cigars during the previous month. Among students in grades 9–12, 30.3% (95% Cl= \pm 2.5%) of those who had smoked cigarettes during the previous month also reported having smoked a cigar, compared with 3.4% (95% Cl= \pm 6.6%) of those who had never smoked a cigarette; among those who had used smokeless tobacco during the previous month, 60.7% (95% Cl= \pm 6.6%) also reported having smoked a cigar during the previous month, compared with 8.3% (95% Cl= \pm 1.0%) of those who had never used smokeless tobacco.

New York Survey

The Roswell Park Cancer Institute survey was conducted in Erie (predominantly urban) and Chautauqua (predominantly rural) counties in New York during the fall of 1996. The survey was administered to 9916 ninth grade students in 57 of the 60 public and parochial high schools in Erie County (81% of the 12,216 ninth grade students in the 60 schools) and to 1677 ninth grade students in 16 of the 18 public schools in Chautauqua County (80% of the 2102 ninth grade students in the 18 schools). Of the students who participated in the survey in Erie County, 79% were non-Hispanic white, 12% were non-Hispanic black, 3% were Hispanic, and 5% were of other racial/ethnic groups. Of those students who participated in the survey in Chautauqua County, 89% were non-Hispanic white. The median age of all students was 14 years. Students

Cigar Smoking Among Teenagers — Continued

completed a self-administered questionnaire with three questions on cigar use and purchasing: "In the past 30 days, did you smoke a cigar?"; "Have you ever bought cigars for yourself?"; and "When you try to buy cigars, how often are you asked about your age?"

Response patterns were similar for the two counties (Table 2). In Erie County, of the 9916 students, 1253 (12.7%) of 9862 students who responded to the question reported having smoked a cigar during the previous 30 days (937 [19.5%] of 4810 boys and 304 [6.1%] of 4983 girls). In Chautauqua County, of the 1677 students, 246 (14.8%) of 1657 students who responded reported having smoked a cigar during the previous

TABLE 2. Number and percentage of ninth grade students who reported having smoked cigars during the previous 30 days or who purchased cigars for their own use, by selected characteristics — Erie and Chautauqua counties, New York, Survey of Alcohol, Tobacco, and Drug Use, 1996

	Eric	e County	*	Chautauqua County [†]			
	Total	Sm or pu	oked rchased	Total	Smoked or purchased		
Category/Characteristic	responses§	No. (%)		responses§	No.	(%)	
SMOKED CIGAR							
Sex							
Male	4810	937	(19.5)	836	201	(24.0)	
Female	4983	304	(6.1)	809	43	(5.3)	
Cigarette use							
Never smoked	6977	323	(4.6)	1147	56	(4.9)	
Occasionally smoked [¶]	1708	458	(26.8)	288	91	(31.6)	
Regularly smoked**	1148	469	(40.9)	218	99	(45.4)	
Smokeless tobacco use							
Not used during previous							
30 days	9469	1032	(10.9)	1532	170	(11.1)	
Used during previous 30 days	348	217	(62.4)	119	75	(63.0)	
Marijuana use							
Never used	6918	360	(5.2)	1126	58	(5.2)	
Ever used	2899	885	(30.5)	521	187	(35.9)	
Used during previous 30 days	1523	606	(39.8)	293	134	(45.7)	
Total	9862	1253	(12.7)	1657	246	(14.8)	
PURCHASED CIGAR							
Sex							
Male	4800	608	(12.7)	831	114	(13.7)	
Female	4969	166	(3.3)	813	21	(2.6)	
Cigarette use							
Never smoked	6957	210	(3.0)	1147	31	(2.7)	
Occasionally smoked [¶]	1705	237	(13.9)	288	35	(12.2)	
Regularly smoked**	1147	331	(28.9)	217	70	(32.3)	
Total	9839	779	(7.9)	1657	136	(8.2)	

* n=9916.

[†]n=1677.

[§]May not equal county totals because of missing data about cigar use and/or purchasing. [¶]Smoked on 1–19 days during the previous 30 days.

**Smoked on 20-30 days during the previous 30 days.

Cigar Smoking Among Teenagers — Continued

30 days (201 [24.0%] of 836 boys and 43 [5.3%] of 809 girls). In comparison, 29.0% of students in Erie County and 30.6% of students in Chautauqua County reported having smoked cigarettes during the previous 30 days. Cigarette smokers also were more likely than noncigarette smokers to report having smoked a cigar during the previous 30 days (Table 2). The prevalence of reported smokeless tobacco use during the previous 30 days was 3.5% in Erie County and 7.3% in Chautauqua County. Among smokeless tobacco users, reported rates of cigar smoking were 62.4% (217 of 348 students who responded) in Erie County and 63.0% (75 of 119 students who responded) in Chautauqua County (Table 2).

Among students who reported ever purchasing a cigar for themselves, most (63.7% in Erie and 77.0% in Chautauqua) also reported having smoked a cigar during the previous 30 days. Among those who had ever purchased a cigar, 76.6% in Erie County and 71.7% in Chautauqua County reported that they were "rarely" or "never" asked about their age when purchasing a cigar. In comparison, 59.0% in Erie County and 67.7% in Chautauqua County reported that they were "rarely" or "never" asked about their age when purchasing a cigar.

Reported by: NJ Kaufman, SL Emont, CR Trimble, CT Orleans, The Robert Wood Johnson Foundation, Princeton, New Jersey. N Briton, T Clark, M Krakow, Health and Addictions Research, Inc, Boston; C Celebucki, D Cullen, G Connolly, Massachusetts Dept of Public Health. A Hyland, J Perla, KM Cummings, Roswell Park Cancer Institute, New York State Dept of Health, Buffalo; A Abdella, K Tippens, Chautauqua County Dept of Health, Mayville, New York. Epidemiology Br, Office on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion, CDC.

Editorial Note: This report is the first to estimate the prevalence of cigar smoking among youth in the United States and documents the level of access to and use of cigars. The risk for several cancers is higher for cigar smokers than for nonsmokers. Therefore, if cigar consumption continues to increase (4), cigar-related morbidity and mortality can be expected to increase.[†]

Although the findings from New York and from Massachusetts were from local surveys, they are consistent with the results from the national survey. However, a potential limitation to these data is that they represent the cigar use of only those adolescents attending school and, therefore, may not be representative of all adolescents.

Although federal law requires states to enact laws prohibiting the sale of cigars and other tobacco products to minors (6), young persons in New York reported being able to purchase cigars easily. These findings, especially if replicated in other communities, may warrant actions to curtail youth access to cigars that are consistent with measures for limiting access to cigarettes and smokeless tobacco (e.g., Food and Drug Administration regulations) (7). The findings from the surveys in this report also indicate that cigar smoking, once primarily an activity among older men (8), is now an activity of both male and female teenagers. Therefore, priorities include the need to further characterize the use of cigars in the United States, determine the prevalence of cigar smoking among adults, and continue monitoring the prevalence of cigar use among youth. Although the Surgeon General's health warning is legally mandated for some tobacco products, the law does not include cigars (9). Therefore, teenagers and other

[†]The National Cancer Institute has announced that it will publish a comprehensive monograph on cigar smoking by the end of 1997 titled "Cigar Smoking in the U.S.: Health Effects and Trends."

Cigar Smoking Among Teenagers — Continued

users of cigars may be unaware of the health risks of cigar smoking. Immediate efforts should be made to publicize the health risks of cigar smoking; deglamorize the product in magazines, movies, and television programs; and protect nonsmokers from secondhand cigar smoke.

References

- US Department of Health and Human Services. The health consequences of smoking: cancer—a report of the Surgeon General. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1982; DHHS publication no. (PHS)82-50179.
- 2. US Department of Health and Human Services. The health consequences of smoking: chronic obstructive lung disease—a report of the Surgeon General. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1984; DHHS publication no. (PHS)84-50205.
- 3. Henningfield JE, Hariharan M, Kozlowski LT. Nicotine content and health risks of cigars. JAMA 1996;276:1857–8.
- US Department of Agriculture. Tobacco situation and outlook report. Washington, DC: US Department of Agriculture, Commodity Economics Division, Economic Research Service, April 1997; document no. TSB-238.
- Roswell Park Cancer Institute. Survey of alcohol, tobacco, and drug use: ninth-grade students in Erie and Chautauqua counties, 1996. Buffalo, New York: Roswell Park Cancer Institute, Department of Cancer Control and Epidemiology, May 1997.
- 6. Substance Abuse and Mental Health Service Administration. Final regulations to implement section 1926 of the Public Health Service Act regarding the sale and distribution of tobacco products to individuals under the age of 18. Federal Register 1996;13:1492–500.
- 7. Food and Drug Administration. Regulations restricting the sale and distribution of cigarettes and smokeless tobacco products to protect children and adolescents: final rule. Federal Register 1996;61:41,314–75.
- Giovino GA, Schooley MW, Zhu B-P, et al. Surveillance for selected tobacco-use behaviors— United States, 1900–1994. MMWR 1994;43(no. SS-3).
- 9. US Department of Health and Human Services. Reducing the health consequences of smoking: 25 years of progress—a report of the Surgeon General. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1989; DHHS publication no. (CDC)89-8411.

Illegal Sales of Cigarettes to Minors — Mexico City, Mexico, 1997

Because of the increasing prevalence of tobacco use among youth in the United States and Mexico (1,2), in 1996 the United States-Mexico Binational Commission (US-MBC) Health Working Group identified prevention of tobacco use, with an emphasis on adolescents, as one of its four priority health concerns. From 1970 to 1990, annual death rates for the leading causes of smoking-related deaths in Mexico nearly tripled and, in 1992, an estimated 10,253 persons in Mexico died as a result of smoking-related diseases, 9% of all deaths that year (3). In addition, from 1988 to 1993, the prevalence of current smoking among minors aged 12–17 years increased from 6.6% to 9.6%, respectively (in Mexico City, the 1993 prevalence was 12.8%), and in 1993, 72% of adult smokers in Mexico reported becoming regular smokers before

Cigarette Sales to Minors — Continued

age 18 years (2,4). Although since 1984 the General Health Law of Mexico has prohibited the sale of tobacco products to minors aged <18 years, compliance with this law has not been assessed. As part of the Mexican national program to reduce the prevalence of cigarette smoking among children and adolescents and in support of the goals of the US-MBC, during 1997 the General Directorate of Epidemiology (GDE) in the Secretariat of Health (SOH) conducted a survey of tobacco outlets in Mexico City to assess the percentage of retailers willing to sell cigarettes to minors. This report summarizes the results of the survey, which indicate that virtually no surveyed retailers asked minors attempting to purchase cigarettes about their age and that most retailers sold cigarettes to minors.

This survey, the first assessment in Mexico of illegal sales of cigarettes to minors, was conducted during March 23–April 4, 1997, in the 16 districts composing Mexico City proper (1990 population: 8.5 million, excluding the surrounding metropolitan area). Because neither commercial business lists of tobacco outlets nor tobacco licensure lists were available and because resources were not available for SOH staff to enumerate a comprehensive list of all operational tobacco outlets in the city, stores were selected as the survey teams visited socioeconomically diverse commercial and residential neighborhoods in each of the 16 districts. Survey teams visited 35 stores in each of 15 districts and 36 stores in one district. The 561 stores included in the non-systematic sample were categorized as small neighborhood stores (302 [54%]), street stalls (137 [24%]), pharmacies (96 [17%]), convenience stores (19 [3%]), and large supermarkets (seven [1%]) (gasoline stations in Mexico are government owned and do not sell cigarettes). Chi-square tests were used to calculate statistical differences in the sales rates associated with selected variables.

The minors who participated in the survey were recruited from the families of staff at GDE and included eight boys aged 10–14 years and seven girls aged 11–15 years. The adult survey escorts were medical residents from the Field Epidemiology Training Program of GDE. Teams consisting of one medical resident, one GDE staff driver, and two minors made one purchase attempt per store using the following protocol: the medical resident entered the store shortly before one of the minors entered the store. The medical resident noted whether age-of-sale warning signs were posted inside the store and unobtrusively observed the transaction between the retailer and the minor as the minor attempted to purchase a pack of cigarettes. If asked by the retailers, the minors were instructed to truthfully state their age and that they carried no age identification. The purchase attempt was considered successful if cigarettes were purchased and was considered unsuccessful if the sale was refused for any reason. If the attempt was successful, the minor promptly left the store with the cigarettes and gave them to the medical resident after the resident exited the store.

Of the 561 stores visited, 443 (79.0%) of the retailers sold cigarettes to the minors (Table 1). Purchase attempts by the oldest minors (aged 14–15 years) were more likely to be successful than those by the youngest minors (aged 10–11 years) (92.2% versus 66.0%, respectively [p<0.01]) and by girls than by boys (84.0% versus 72.7%, respectively [p<0.01]). Sales were transacted at all types of stores. Although the proportion of successful sales did not vary by sex of the retailer, the proportion was higher for attempts involving male clerks and girls than for those involving male clerks and boys (88.3% versus 68.1%, p<0.01). Age-of-sale warning signs were displayed in 64 (11.8%) stores; the presence of a warning sign was not associated with lower sales rates. Four

Cigarette Sales to Minors — Continued

	No retail	S	uccessful attem	pts
Category	businesses	No.	(%)	p value
Age group (yrs) of minor				
10–11	247	163	(66.0)	<0.01
12–13	44	31	(70.5)	
14–15	270	249	(92.2)	
Sex of minor				
Male	253	184	(72.7)	<0.01
Female	308	259	(84.0)	
Type of store				
Large supermarket	7	3	(42.9)	<0.01
Convenience	19	11	(57.9)	
Small neighborhood	302	238	(78.8)	
Pharmacy	96	71	(74.0)	
Street stalls	137	120	(87.6)	
Warning sign				
Yes	64	47	(73.4)	0.25
No	497	396	(79.7)	
Sex of retailer				
Male	300	237	(79.0)	0.98
Female	261	206	(78.9)	
Total	561	443	(79.0)	

TABLE 1. Number of retail businesses surveyed and number and percentage of
successful attempts by minors* to purchase cigarettes, by category - Mexico City,
Mexico, 1997

*Persons aged <18 years.

(0.7%) retailers asked the minor's age; one (0.2%) asked for proof of age; and 30 (5.4%) asked for whom the cigarettes were being purchased. Of the 118 retailers who did not sell cigarettes to the participating minors, 73 (62%) indicated to the participants that they do not sell cigarettes to minors.

Reported by: P Kuri-Morales, MD, P Cravioto, MSc, M Hoy, MD, S Huerta, MD, A Revuelta, MD, B Jasso, MD, R Tapia-Conyer, MD, Secretariat of Health, Mexico City, Mexico. Office on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion, CDC.

Editorial Note: Most of the retailers included in the sample in this survey in Mexico City illegally sold cigarettes to the participating minors. In the United States, a national health objective for 2000 is to reduce to $\leq 20\%$ the proportion of retailers who sell tobacco to minors (objective 3.13) (5). Among 13 local U.S. studies published during 1989–1993, rates of over-the-counter cigarette sales to minors ranged from 32% to 87% (6). Compliance surveys estimating the overall rate of cigarette sales to minors also have been conducted in Canada (52.1% in 1995 and 39.5% in 1996) (7) and Adelaide, Australia (46% in 1991) (8).

The findings in this report are subject to at least two limitations. First, because this survey used a nonsystematic sample of retail businesses, the findings probably do not uniformly represent the patterns of tobacco sales to minors throughout Mexico City. For example, even though the survey teams visited all districts of Mexico City, some types of stores and neighborhoods at some socioeconomic levels—especially those at

Cigarette Sales to Minors - Continued

lower levels—may not have been included in the sample. However, it is not known whether sales rates in lower socioeconomic neighborhoods differed from those in higher socioeconomic neighborhoods. Second, the rate may have been underestimated because retailers in small neighborhood stores and street stalls in particular may have suspected that the adult team member, who entered the store or approached the stall before the minor, was accompanying the minor.

Based on current global patterns of smoking, the World Health Organization (WHO) has projected that 200-300 million persons who are aged <20 years in 1997 will die from smoking-related diseases later in life (9). In 1986, the World Health Assembly adopted a resolution urging member states to consider a comprehensive tobaccocontrol strategy containing nine elements (10), including one that targets the prevention of smoking by children and adolescents. However, in the early 1990s, WHO determined that only approximately 25 countries had established laws prohibiting the sale of cigarettes to minors (the age of prohibition varied from 16 to 21 years), and that among these, only a limited number had attempted to enforce the laws. To decrease cigarette sales to minors, WHO recommends that countries adopt the following four measures: 1) establish a minimum age of purchase of 18 years or older; 2) create a tobacco-sales licensing system to identify tobacco retailers and inform them of their legal responsibilities; 3) establish a graduated schedule of civil law penalties for illegal sales, ranging from warnings to license revocations; and 4) enlist the assistance of teenagers in efforts of enforcement officers to assess retailers' compliance with the prohibition of sale to minors. Other categories of legislation also may be effective in decreasing sales to minors. For example, several local studies in the United States demonstrated substantially reduced tobacco sales to minors when retailers requested photo identification or other proof of age from persons attempting to purchase tobacco products (1).

SOH will use the results of this survey to emphasize the need for assessing compliance of retailers in other cities with the federal law prohibiting tobacco sales to minors in Mexico and to underscore the need for resources to support increased enforcement activities. In addition to the enforcement of strong minors' access laws, a comprehensive approach for preventing initiation of smoking by youth should include provisions that reduce the appeal of cigarettes to minors through restrictions on advertising and promotion and through educational programs (1).

References

- 1. Food and Drug Administration. Regulations restricting the sale and distribution of cigarettes and smokeless tobacco products to protect children and adolescents: final rule. Federal Register 1996;61:41,314–75.
- 2. Secretariat of Health. The National Addiction Survey [Spanish]. Mexico City, Mexico: Secretariat of Health, General Directorate of Epidemiology, 1993.
- 3. CDC. Smoking-attributable mortality—Mexico, 1992. MMWR 1995;44:372-3,379-81.
- 4. Tapia-Conyer R, Medina-Mora ME, Sepulveda J, et al. The National Addiction Survey of Mexico [Spanish]. Salud Pública Mex 1990;32:507–22.
- Public Health Service. Healthy people 2000: national health promotion and disease prevention objectives—midcourse review and 1995 revisions. Washington, DC: US Department of Health and Human Services, Public Health Service, 1995.
- 6. US Department of Health and Human Services. Preventing tobacco use among young people: a report of the Surgeon General. Atlanta, Georgia: US Department of Health and Human Services, Public Health Service, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1994.

Cigarette Sales to Minors — Continued

- 7. AC Nielsen. Measurement of retailer compliance with respect to tobacco sales to minors legislation: wave 2, 1996. Montreal, Quebec: AC Nielsen, 1996.
- 8. Wakefield M, Carrangis J, Wilson D, Reynolds C. Illegal cigarette sales to children in South Australia. Tobacco Control 1992;1:114–7.
- 9. World Health Organization. Guidelines for controlling and monitoring the tobacco epidemic. Geneva, Switzerland: World Health Organization, 1997 (in press).
- World Health Organization. Tobacco or health: first global status report. Geneva, Switzerland: World Health Organization, 1997 (in press).

As part of its commemoration of CDC's 50th anniversary, MMWR is reprinting selected MMWR articles of historical interest to public health, accompanied by a current editorial note. Reprinted below is the report published October 30, 1987, which analyzed smoking-attributable mortality and years of potential life lost for 1984, followed by a contemporary editorial note.

Perspectives in Disease Prevention and Health Promotion

Smoking-Attributable Mortality and Years of Potential Life Lost — United States, 1984

Cigarette smoking has been identified as the chief avoidable cause of death in the United States (1). Several estimates of mortality attributable to cigarette smoking have been reported, including 270,000 deaths for 1980 (2) and 314,000 deaths for 1982 (3). Published estimates vary considerably because of changing mortality rates, decreasing smoking rates, and differences in methods used. Smoking-attributable mortality and years of potential life lost (YPLL) for 1984 are analyzed in this report.

Relative risk (RR) estimates for smoking-related diseases and prevalence estimates of current, former, and never smokers among adults \geq 20 years of age were used to calculate the smoking-attributable fraction (SAF) and smoking-attributable mortality for 19 underlying causes of death (2) (Table 1).* Age-, sex-, and race-specific mortality data for 1984 were obtained from National Center for Health Statistics reports. Age-, sex-, and race-specific smoking prevalence rates were obtained from the 1985 Current Population Survey (Supplement) of the Bureau of the Census (Office on Smoking and Health, CDC, unpublished data). Years of potential life lost were calculated to age 65 according to previously described methods (6). Age-adjusted smoking-attributable mortality and YPLL rates were calculated by the direct method, with the 1984 U.S. population used as the standard.

For deaths among adults, the disease-specific SAFs are derived from RR estimates for current and former smokers that are weighted averages from four prospective studies (7–10). RR estimates for women based on these studies may be lower than the current RRs for many of the specific smoking-related diseases among women. However, the SAF for lung cancer among women (0.75) has been updated based on RR

^{*}The equation for calculating the smoking-attributable fraction of each disease category is: SAF=[p₀ + p₁(RR₁) + p₂(RR₂)] - 1/[p₀ + p₁(RR₁) + p₂(RR₂)] where p₀=percentage of never smokers, p₁=percentage of current smokers, p₂=percentage of former smokers, RR₁=relative risk for current smokers (relative to never smokers), and RR₂=relative risk for former smokers (relative to never smokers) (*4*). This formula is derived from the standard attributable risk (AR) formula (*5*): AR=p(RR - 1)/[p(RR - 1) + 1].

Smoking-Attributable Mortality — Continued

estimates from more recent mortality data (11). Race-specific RR estimates for smoking-attributable diseases were not available.

For four pediatric diagnoses, the mortality attributed to maternal smoking during pregnancy for children <1 year of age was determined. These calculations used RR estimates from McIntosh (12) and current smoking prevalence among women

TABLE 1. Total mortality, weighted	I smoking-attributable fractions (SAF), and
smoking-attributable mortality (SAM)	, by disease category and sex — United States,
1984	

	_	Males			Total			
Disease Ca	tegory*	Deaths	SAF	SAM	Deaths	SAF	SAM	SAM [†]
Adults ≥20	years old							
Neoplasms	5:							
140-149	Lip, oral cavity, pharynx	5,754	0.688	3,958	2,689	0.413	1,110	5,068
150	Esophagus	6,310	0.589	3,717	2,345	0.536	1,257	4,974
151	Stomach	8,468	0.172	1,455	5,772	0.254	1,467	2,922
					11,634			
157	Pancreas	11,513	0.300	3,459		0.142	1,653	5,112
161	Larynx	2,959	0.806	2,385	664	0.413	274	2,660
162	Trachea, lung, bronchus	82,459	0.796	65,659	36,227	0.750	27,170	92,829
180	Cervix uteri	0	0.0	0	4,562	0.369	1,685	1,685
188	Urinary bladder	6,597	0.371	2,447	3,114	0.274	853	3,299
189	Kidney, other urinary	5,424	0.243	1,319	3,403	0.118	403	1,722
Circulatory	diseases:							
401-405	Hypertension	13,464	0.156	2,099	17,855	0.148	2,645	4,744
410-414	lschemic heart disease <age 65<="" td=""><td>78,340</td><td>0.285</td><td>22,362</td><td>27,000</td><td>0.181</td><td>4,892</td><td>27,253</td></age>	78,340	0.285	22,362	27,000	0.181	4,892	27,253
410-414	lschemic heart disease ≥age 65	211,003	0.159	33,461	224,756	0.075	16,816	50,276
427.5	Cardiac arrest	19,392	0.399	7,745	17,296	0.344	5,950	13,695
430-438	Cerebrovascular disease	59,185	0.096	5,692	88,285	0.139	12,228	17,920
440	Arteriosclerosis	9,235	0.238	2,200	15,216	0.315	4,797	6,996
441	Aortic aneurysm	10,323	0.624	6,444	4,791	0.468	2,244	8,689
Respiratory	/ Diseases:							
480-487	Pneumonia, influenza	28,774	0.208	5,986	28,935	0.093	2,679	8,664
401 402	emphysema	10,708	0.850	9,097	5,517	0.694	3,831	12,928
496	Chronic airways obstruction	31,240	0.850	26,541	16,625	0.694	11,545	38,085
Digestive d	liseases:							
531-534	Ulcers	3,251	0.479	1,556	3,365	0.445	1,497	3,053
Pediatric di	iseases, <1 year old							
765	Short gestation, low birthweight	1,729	0.182	314	33	0.182	279	593
769	Respiratory distress syndrome	2,178	0.182	396	1,379	0.182	251	647
770	Other respiratory conditions							
	of newborn	1,982	0.182	360	1,515	0.182	275	636
798.0	Sudden infant death syndrome	3,176	0.128	405	2,069	0.128	264	669
Total [†]				209,057			106,063	315,120

* International Classification of Disease, ninth revision. [†] Sums may not equal total because of rounding.

Smoking-Attributable Mortality — Continued

20–64 years of age as a proxy for the percentage of pregnant women who smoke. The RR (1.50) for sudden infant death syndrome from McIntosh (*12*) was used, but the RR (1.76) for total infant mortality reported by McIntosh was used to calculate the SAF for only three specific infant death categories (short gestation/low birthweight, respiratory distress syndrome, and other respiratory conditions).

An estimated 315,120 deaths and 949,924 YPLL before age 65 years resulted from cigarette smoking in 1984 (Table 2). The smoking-attributable mortality rate among men is more than twice the rate among women, and the rate among blacks is 20% higher than the rate among whites (Table 3). The smoking-attributable YPLL rate among men is more than twice the rate among women, and the rate among blacks is more than twice the rate among whites (Table 3).

Reported by: Office on Smoking and Health, Center for Health Promotion and Education, CDC. **Editorial Note**: The total smoking-attributable mortality and YPLL reported here is similar to that cited in previous reports (2,3), showing that the disease impact of smoking in the United States continues to be enormous despite recent declines in the prevalence of smoking. These figures do not include mortality and YPLL due to peripheral vascular disease (for which specific RR estimates are generally lacking), cancer at unspecified sites, cigarette-caused fires, or involuntary (passive) smoking. In 1984, an estimated 1,570 deaths were attributed to cigarette-initiated fires (13); an estimated 3,825 nonsmokers per year die from lung cancer attributed to involuntary smoking are included, the estimated total of smoking-attributable deaths in the United States in 1984 is 320,515, or 15.7% of all (2,039,369) U.S. deaths. Total smoking-attributable YPLL (949,924) represents

		Mortality			YPLL	
	Males	Females	Total [†]	Males	Females	Total [†]
Whites	184,296	95,340	279,636	489,827	199,590	689,418
Blacks	22,647	10,131	32,779	129,952	63,473	193,425
Total population [§]	209,057	106,063	315,120	661,651	288,273	949,924

TABLE 2. Estimated smoking-attributable mortality and years of potential life lost (YPLL)*, by race and sex — United States, 1984

*YPLL before age 65.

[†]Sums may not equal total because of rounding.

§Includes whites, blacks, and racial category "other."

TABLE 3. Age-adjusted	smoking-attributable	mortality rates*	and years of	of potential
life lost (YPLL) rates [†] , by	/ race and sex — Unit	ed States, 1984		

		Mortality rate)			
	Males	Females	Total [†]	Males	Females	Total
Whites	189.7	64.2	119.0	5.56	2.17	3.81
Blacks	236.5	75.5	143.2	12.07	4.85	8.14
Total population [§]	192.6	68.0	133.2	6.53	2.71	4.56

*Per 100,000 persons (population data from 1984 U.S. Census).

[†]YPLL before age 65/1,000 persons <65 years (population data from 1984 U.S. Census).

§Includes whites, blacks, and racial category "other."

Smoking-Attributable Mortality — Continued

8.1% of all (11,761,000) U.S. YPLL before age 65 (excluding YPLL due to cigarettecaused fires or involuntary smoking).

Among blacks, the smoking-attributable mortality (32,779) represents 13.9% of total 1984 mortality (235,884), whereas the smoking-attributable mortality for whites (279,636) was 15.7% of total 1984 mortality (1,781,897), excluding deaths due to fires or involuntary smoking. However, the smoking-attributable mortality rate and YPLL rate were higher among blacks than among whites. These differences in rates reflect a higher prevalence of smoking and a higher mortality rate from smoking-related diseases among blacks. Higher YPLL rates among blacks may also reflect more smokingattributable deaths at earlier ages. Because blacks tend to smoke fewer cigarettes per day than whites (*15,16*), the difference in smoking-attributable mortality and YPLL rates between blacks and whites may be slightly overestimated. On the other hand, the RR of smoking-related diseases among blacks may be higher than the RR estimates used here because of increased interactions between smoking and other risk factors, different tar and nicotine exposures, or different smoking patterns. Still, these findings support previously cited concerns regarding the increased burden of smoking-related disease among blacks (*17*).

Smoking prevalence for 1985 was used to calculate the SAFs in this study. However, the 1984 smoking-related mortality is a result of a higher smoking prevalence during the 1950s, '60s, and '70s, the decades during which these diseases were developing. Therefore, the SAFs used here are conservative.

CDC has examined YPLL before age 65 years since 1979 (6). In this study, most smoking-related deaths (218,691, or 69.4%) occurred among persons \geq 65 years of age. Thus, the smoking-attributable YPLL among persons <65 reported here (949,924) is substantially lower than the 3.6 million smoking-attributable YPLL calculated when the average life expectancy in the United States is used for calculating YPLL for 1984.

Group-specific calculations such as these are possible for states and other defined populations if mortality and smoking prevalence data for those populations are available. A computer program has recently been developed to aid in calculating mortality and YPLL attributed to cigarette smoking (*18*). CDC is now collaborating with all 50 state health departments, Puerto Rico, and the District of Columbia to perform similar studies. Results from this project will be reported in 1988.

References

- Office on Smoking and Health. The health consequences of smoking: cancer—a report of the Surgeon General. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, 1982:xi; DHHS publication no. (PHS)82-50179.
- 2. Rice DP, Hodgson TA, Sinsheimer P, Browner W, Kopstein AN. The economic costs of the health effects of smoking, 1984. Milbank Mem Fund Q 1986;64:489-547.
- 3. Office of Technology Assessment. Smoking-related deaths and financial costs. OTA Staff Memorandum. Health Program, U.S. Congress, 1985.
- 4. Walter SD. The estimation and interpretation of attributable risk in health research. Biometrics 1976;32:829-49.
- Lilienfeld AM, Lilienfeld DE. Foundations of epidemiology. 2nd ed. New York: Oxford University Press, 1980.
- CDC. Premature mortality in the United States: public health issues in the use of years of potential life lost. MMWR 1986;35(suppl 2S).
- Hammond EC. Smoking in relation to the death rates of one million men and women. In: Haenszel WM, ed. Epidemiological approaches to the study of cancer and other chronic diseases. Bethesda: National Cancer Institute, US Department of Health, Education, and Welfare, Public Health Service, 1966:127-204. (NCI Monograph no. 19).

Smoking-Attributable Mortality — Continued

- 8. Doll R, Peto R. Mortality in relation to smoking: 20 years' observations on male British Doctors. Br Med J 1976;2:1525-36.
- 9. Doll R, Gray R, Hafner B, Peto R. Mortality in relation to smoking: 22 years' observations on female British doctors. Br Med J 1980;280:967-71.
- Cederlöf R, Friberg L, Lundman T. The interactions of smoking, environment, and heredity and their implications for disease etiology: a report of epidemiological studies on the Swedish twin registries. Acta Med Scand 1977;612(suppl):7-128.
- 11. American Cancer Society. 1986 cancer facts and figures. New York: American Cancer Society, 1986:17.
- 12. McIntosh ID. Smoking and pregnancy: attributable risks and public health implications. Can J Public Health 1984;75:141-8.
- Hall JR Jr. Expected changes in fire damages from reducing cigarette ignition propensity. Report No. 5, Technical Study Group, Cigarette Safety Act of 1984. Quincy, Massachusetts: National Fire Protection Association, Fire Analysis Division, 1987.
- 14. National Academy of Sciences. Environmental tobacco smoke: measuring exposures and assessing health effects. Washington, DC: National Academy Press, 1986: Appendix D.
- 15. CDC. Cigarette smoking in the United States, 1986. MMWR 1987;36:581-5.
- National Center for Health Statistics. Health, United States, 1986. Washington, DC: US Department of Health and Human Services, Public Health Service, 1986:126; DHHS publication no. (PHS)87-1232.
- 17. CDC. Cigarette smoking among blacks and other minority populations. MMWR 1987;36:404-7.
- Shultz JM. SAMMEC: smoking-attributable mortality, morbidity, and economic costs (computer software and documentation). Minnesota: Center for Nonsmoking and Health, Minnesota Department of Health, 1986.

Editorial Note—1997: In 1987, CDC published the preceding report that provided a detailed and comprehensive estimate of the number of deaths attributed to cigarette smoking in the United States. Using the attributable fraction, which measures the magnitude of a public health problem accounted for by an etiologic agent, CDC was able to quantify the impact of smoking. This method established that smoking was the leading cause of preventable deaths in the United States (1). As a result, increased emphasis was placed on decreasing the health burden caused by tobacco use and on reducing cigarette smoking. Since this SAM estimate was published in 1987, continued research has increased understanding of the health risks associated with tobacco use, including nicotine addiction and the recognition that addiction begins in childhood. Public health programs have responded by focusing on preventing tobacco use among adolescents, assisting in tobacco-use cessation, and protecting nonsmokers from environmental tobacco smoke. This contemporary editorial note reviews previous SAM estimates, presents new SAM estimates for 1990–1994, and discusses future implications.

SAM and YPLL estimates for the United States published since the first estimate for 1984 include 390,000 deaths for 1985, 434,000 deaths and 6 million YPLL before age 85 for 1988, and 418,000 deaths and 5 million YPLL to life expectancy for 1990 (*2*). SAM and YPLL also have been estimated for all 50 states and the District of Columbia for 1985 and for 1990 (*3*). Although all estimates were calculated by the same equation used for the SAF, the data sources, study populations, and causes of death have changed. The Smoking-Attributable Mortality, Morbidity, and Economic Costs (SAM-MEC) software program has also been used for calculating these estimates (*2*).

Since 1989, RR estimates for calculating SAM and YPLL have been obtained from the American Cancer Society's Cancer Prevention Study II (CPS-II) for 1982–1986 (4). The CPS-II was selected, in part, because it is the largest prospective U.S. study that has collected data on the relation between smoking and mortality (4). Recent SAM

Smoking-Attributable Mortality — Continued

estimates for adults have been limited to persons aged \geq 35 years because the CPS-II study population was restricted to this age range. Deaths from stomach cancer and ulcers were dropped from the calculation of SAM because a causal relation has not been established (4). Conversely, the cardiovascular and respiratory disease categories were expanded to include the *International Classification of Diseases, Ninth Revision* [ICD-9], codes 390–398, 415–417, 420–429, 442–448, 010–012, and 493. The CPS-II data also enabled the calculation of the RR for smoking and cerebrovascular disease, which declines with age (4), for two age groups (35–64 years and \geq 65 years).

Cigarette smoking remains the leading preventable cause of death in the United States. The same methods and data sources that were used to calculate the 1990 SAM and YPLL (2)[†] were used for the 1990–1994 calculations, which indicated that 2,153,700 deaths (1,393,200 men and 760,400 women; total annual average: 430,700 deaths) were attributed to smoking (19.5% of all deaths). A total of 906,600 of these deaths resulted from cardiovascular diseases; 778,700, from neoplasms; 454,800, from nonmalignant respiratory diseases; 7900, from diseases among infants; and 5500, from smoking-related fires. Lung cancer (616,800 deaths), ischemic heart disease (IHD) (490,000 deaths), and chronic airway obstruction (270,100 deaths) accounted for most deaths. During 1990–1994, cigarette smoking resulted in 5,732,900 YPLL before age 65 years and in 28,606,000 YPLL to life expectancy.

During 1990–1994, estimates of SAM were higher among men than among women, reflecting their longer duration and higher prevalence of smoking and greater numbers of cigarettes smoked per day (6). Annual SAM rates will probably remain stable if current trends in smoking prevalence among adults continue. Although the prevalence of smoking among persons aged \geq 35 years decreased from 1985 to 1990 (28.4% to 24.1%), during 1990–1994, smoking prevalence remained relatively constant—at 23.6%–24.8% (CDC, unpublished data). However, the prevalence of smoking patterns continue into adulthood, SAM and YPLL are expected to increase. Assuming that one third of adult smokers, 10% of former smokers, and 5.3 million persons aged <18 years die from smoking and that current smoking patterns continue, an estimated 25 million persons alive today will die prematurely from smoking-related illnesses (7,8).

Lung cancer has been and probably will continue to be the leading cause of SAM because, although lung cancer death rates are decreasing among men, rates are continuing to increase among women (9). Among women, death rates for lung cancer have surpassed those for breast cancer since 1987 (9). In addition, because recent trends indicate a slowing of the decline in IHD mortality, IHD will probably remain a major contributor to SAM (9).

SAM and YPLL may be underestimated for several reasons (2); recent studies have addressed two of these reasons. First, SAM and YPLL estimates are based on the prevalence of current and former smokers in the current year; however, the deaths that occur during a given year are primarily among persons who began smoking 30–50 years earlier (10), many of whom have quit smoking (10). Including these persons in the prevalence estimates of former smokers may decrease the SAF because the summary measure of risk for former smokers does not reflect their increased likeli-

⁺Except for the prevalence of smoking among pregnant women in the United States for 1992 through 1994, which was estimated from the 1992–1993 National Pregnancy and Health Survey (5).

Smoking-Attributable Mortality — Continued

hood of dying from a smoking-related disease (4). Among whites, expanding the classification of smoking to include information on duration and number of cigarettes smoked per day resulted in 10% larger SAM estimates for IHD than SAM estimates in which smoking was categorized as current, former, and never (10). Second, the SAM estimates do not include mortality caused by cigar smoking, pipe smoking, or smokeless tobacco use. Approximately 1000 deaths were attributable to pipe smoking in 1991 (11).

Although SAM and YPLL estimates are not adjusted for confounders (2–4), a recent study has documented little change in SAM estimates after adjustment for confounders (12). Among whites, SAM estimates for the combined disease categories of lung cancer, IHD, bronchitis/emphysema, chronic airway obstruction, and cerebrovascular disease were 2% higher than age-adjusted estimates after adjustment for relevant confounders including age, education, alcohol intake, diabetes, and hypertension (12).

Cigarette smoking has resulted in approximately 10 million deaths since the first Surgeon General's report on smoking and health in 1964 (*2,4,13*). In 1993, \$50 billion in medical costs were attributable to smoking (*14*). The human and economic costs of smoking will continue to accumulate until the completely effective implementation of public health efforts to prevent initiation, to promote cessation, and to protect nonsmokers from the adverse effects of environmental tobacco smoke. Examples of such efforts include Food and Drug Administration regulations to restrict youth access to tobacco and to reduce the appeal of cigarette advertising to youth (*7*); comprehensive state-based efforts, including tax increases and earmarked funding for tobacco-use prevention and mass media campaigns similar to those in Massachusetts and California (*15*); physician adherence to the Agency for Health Care Policy and Research's smoking cessation guidelines (*8*); institutional adoption of the Guidelines for School Health Programs to Prevent Tobacco Use and Addiction (*16*); and clean indoor-air policies that protect nonsmokers.

1997 Editorial Note by Ann M Malarcher, PhD, Jeffrey H Chrismon, Gary A Giovino, PhD, Michael P Eriksen, ScD, Office on Smoking and Health, National Center for Chronic Disease Prevention and Health Promotion, CDC.

References

- 1. McGinnis JM, Foege WH. Actual causes of death in the United States. JAMA 1993;270:2207–12.
- 2. CDC. Cigarette smoking-attributable mortality and years of potential life lost—United States, 1990. MMWR 1993;42:645–9.
- Nelson DE, Kirkendall RS, Lawton RL, et al. Surveillance for smoking-attributable mortality and years of potential life lost, by state—United States, 1990. MMWR 1994;43(no. SS-1):1–8.
- 4. US Department of Health and Human Services. Reducing the health consequences of smoking: 25 years of progress—a report of the Surgeon General. Washington, DC: US Department of Health and Human Services, Public Health Service, CDC, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1989.
- National Institute on Drug Abuse. Summary tables: annualized estimates from the National Pregnancy and Health Survey. Rockville, Maryland: US Department of Health and Human Services, Public Health Service, National Institutes of Health, National Institute on Drug Abuse, 1994.
- Giovino GA, Schooley MW, Zhu B-P, et al. Surveillance for selected tobacco-use behaviors— United States, 1900–1994. MMWR 1994;43(no. SS-3).
- 7. CDC. Projected smoking-related deaths among youth—United States. MMWR 1996;45:971-4.
- 8. CDC. State specific prevalence of cigarette smoking—United States, 1995. MMWR 1996;45:962–6.
- National Center for Health Statistics. Health, United States, 1995. Hyattsville, Maryland: US Department of Health and Human Services, Public Health Service, CDC, 1996.

Smoking-Attributable Mortality — Continued

- Schulman J, Mowery PD, Pierce BK, et al. Methodologic issues in estimating smoking attributable mortality. Atlanta, Georgia: Battelle, Centers for Public Health Research and Evaluation, 1995.
- 11. Nelson DE, Davis RM, Chrismon JH, Giovino GA. Pipe smoking in the United States, 1965–1991: prevalence and attributable mortality. Prev Med 1996;25:91–9.
- Schulman J, Epstein L, Mowery PD, Pierce B, Euskirchen E, Abed J. Smoking attributable mortality: control for confounding. Atlanta, Georgia: Battelle, Centers for Public Health Research and Evaluation, 1997.
- 13. US Department of Health, Education, and Welfare. Smoking and health report of the advisory committee to the Surgeon General of the Public Health Service. Washington, DC: US Department of Health, Education, and Welfare, Public Health Service, 1964.
- 14. CDC. Medical-care expenditures attributable to cigarette smoking—United States, 1993. MMWR 1994;43:469–72.
- 15. CDC. Cigarette smoking before and after an excise tax increase and an antismoking campaign—Massachusetts, 1990–1996. MMWR 1996;45:966–70.
- 16. CDC. Guidelines for school health programs to prevent tobacco use and addiction. MMWR 1994;43(no. RR-2).

Outbreaks of Cyclosporiasis — United States, 1997

In April and May 1997, CDC received reports of seven event-associated clusters of cases of cyclosporiasis from five states (California, Florida, Nevada, New York, and Texas). Approximately 80 cases of infection with human-associated *Cyclospora*, a recently characterized coccidian parasite (1), have been laboratory-confirmed. State and local health departments, CDC, and the Food and Drug Administration are conducting investigations to identify the vehicles of infection.

Both foodborne and waterborne outbreaks of cyclosporiasis have previously been reported in the United States during spring and summer months (2-4). In 1996, a total of 978 laboratory-confirmed cases of cyclosporiasis in the United States and Canada were reported in association with a widespread foodborne outbreak (3). The average incubation period of cyclosporiasis is 1 week. Illness can be protracted (from days to weeks) with frequent, watery stools and other gastrointestinal symptoms; symptoms may remit and relapse. Health-care providers should consider *Cyclospora* infection in persons with prolonged diarrheal illness and specifically request laboratory testing for this parasite (5, 6), which is not routinely performed by most laboratories.

Cyclosporiasis can be treated with a 7-day course of oral trimethoprim (TMP)sulfamethoxazole (SMX) (for adults, 160 mg TMP plus 800 mg SMX twice daily; for children, 5 mg/kg TMP plus 25 mg/kg SMX twice daily) (7). Treatment regimens for patients who cannot tolerate sulfa drugs have not yet been identified.

Health departments that identify cases of cyclosporiasis should contact CDC's Division of Parasitic Diseases, National Center for Infectious Diseases, telephone (770) 488-7760.

Reported by: State and local health departments. Office of Regulatory Affairs, and Center for Food Safety and Applied Nutrition, Food and Drug Administration. Div of Parasitic Diseases, National Center for Infectious Diseases, CDC.

References

- 1. Ortega YR, Sterling CR, Gilman RH, Cama VA, Díaz F. Cyclospora species—a new protozoan pathogen of humans. N Engl J Med 1993;328:1308–12.
- 2. CDC. Update: outbreaks of *Cyclospora cayetanensis* infection—United States and Canada, 1996. MMWR 1996;45:611–2.

452

Cyclosporiasis — Continued

- 3. Herwaldt BL, Ackers M-L, Cyclospora Working Group. An outbreak in 1996 of cyclosporiasis associated with imported raspberries. N Engl J Med 1997;336:1548–56 (in press).
- 4. Huang P, Weber JT, Sosin DM, et al. The first reported outbreak of diarrheal illness associated with *Cyclospora* in the United States. Ann Intern Med 1995;123:409–14.
- 5. Soave R. Cyclospora: an overview. Clin Infect Dis 1996;23:429-37.
- 6. Garcia LS, Bruckner DA. Diagnostic medical parasitology. 3rd ed. Washington, DC: American Society for Microbiology, 1997:66–9.
- 7. Hoge CW, Shlim DR, Ghimire M, et al. Placebo-controlled trial of co-trimoxazole for Cyclospora infections among travellers and foreign residents in Nepal. Lancet 1995;345:691–3.

Notice to Reader

Courses on Epidemiology, Public Health Informatics, and Advanced Case Studies in Public Health Practice

CDC and the University of Washington School of Public Health and Community Medicine will cosponsor courses in epidemiology, public health informatics, and advanced case studies in public health practice as part of the 1997 Summer Institute of Public Health Practice at the University of Washington School of Public Health, July 21–August 1, 1997, in Seattle, Washington. The epidemiology course will provide an in-depth consideration of special epidemiologic topics. The informatics course will cover the principles of applying information technology to public health practice. The advanced case studies course will examine current issues in public health practice using several management tools and analytic techniques. These courses are offered in conjunction with various practice-oriented public health science and management courses at the Summer Institute. Public health practitioners are encouraged to attend the courses.

Participation in each course is limited, and early application is encouraged. There is a tuition charge. Additional information and applications are available from the Center for Health Education and Research, 1001 Broadway, Suite 217, Seattle, WA 98122; telephone (206) 720-4250; fax (206) 720-4218; e-mail: si97@u.washington.edu; World-Wide Web site: http://healthlinks.washington.edu/nwcphp/edu/si/intro.html.



FIGURE I. Selected notifiable disease reports, comparison of provisional 4-week totals ending May 17, 1997, with historical data — United States

*Ratio of current 4-week total to mean of 15 4-week totals (from previous, comparable, and subsequent 4-week periods for the past 5 years). The point where the hatched area begins is based on the mean and two standard deviations of these 4-week totals.

TABLE I. Summary — provisional cases of selected notifiable diseases, United States, cumulative, week ending May 17, 1997 (20th Week)

	Cum. 1997		Cum. 1997
Anthrax Brucellosis Cholera Congenital rubella syndrome Cryptosporidiosis* Diphtheria Encephalitis: California* eastern equine* St. Louis* western equine* Hansen Disease Hantavirus pulmonary syndrome*† Hemolytic uremic syndrome, post-diarrheal* HIV infection, pediatric* [§]	16 1 2 430 4 4 1 1 40 4 4 14 92	Plague Poliomyelitis, paralytic Psittacosis Rabies, human Rocky Mountain spotted fever (RMSF) Streptococcal disease, invasive Group A Streptococcal toxic-shock syndrome* Syphilis, congenital [¶] Tetanus Toxic-shock syndrome Trichinosis Typhoid fever Yellow fever	1 15 2 49 542 13 62 9 41 3 102

-:no reported cases

*Not notifiable in all states. [†]Updated weekly from reports to the Division of Viral and Rickettsial Diseases, National Center for Infectious Diseases (NCID). ³Updated monthly to the Division of HIV/AIDS Prevention–Surveillance and Epidemiology, National Center for HIV, STD, and TB Prevention (NCHSTP), last update April 29, 1997. ¹Updated from reports to the Division of STD Prevention, NCHSTP.

					Esche	erichia						
	All	AIDS		nydia	COII O	157:H7 PHLIS [§]	Gono	rrhea	C/NA,NB			
Reporting Area	Cum. 1997*	Cum. 1996	Cum. 1997	Cum. 1996	Cum. 1997	Cum. 1997	Cum. 1997	Cum. 1996	Cum. 1997	Cum. 1996		
UNITED STATES	20,222	26,411	145,458	155,853	374	166	92,582	115,544	1,122	1,235		
NEW ENGLAND	671	942	6,130	7,247	33	15	2,072	3,016	17	41		
Maine N.H.	25 8	15 31	333	270	1	-	17 50	18 56	- 3	- 2		
Vt.	16	9	146	175	2	1	20	24	-	12		
R.I.	282 55	549 61	2,703	2,542 808	24	- 14	193	832 208	2	24		
Conn.	285	277	1,871	3,452	3	-	931	1,878	-	-		
MID. ATLANTIC	6,683 1 143	7,513 794	17,953 N	27,522 N	27 16	4	10,483	16,606 2 725	126 98	104 83		
N.Y. City	3,308	4,474	9,220	14,659	5	-	4,219	6,437	-	2		
N.J. Pa	1,444 788	1,413 832	2,742 5,991	5,675 7,188	6 N	- 1	1,584 2,879	3,513 3,931	- 28	- 19		
E.N. CENTRAL	1,416	2,208	22,901	34,445	65	24	13,848	22,231	224	201		
Ohio	270	487	5,326	7,804	20	10	3,391	5,540	5	4		
III.	302 509	980	3,204 4,357	3,891 9,965	13	5	2,205 2,123	6,593	16	48		
Mich.	259	318	7,309	8,514	17 N	2	4,950	5,671	198	143		
WIN CENTRAL	283	570	2,705	4,271	50	33	3 955	5 601	- 60	- 22		
Minn.	79	125	U,424	1,806	29	20	U,555	867	1	-		
lowa Mo	59 150	43 232	1,765 4,168	1,503 5.325	12	5 5	452 2.766	398 3.201	15 31	8		
N. Dak.	4	5	339	404	3	2	23	11	2	-		
S. Dak. Nebr.	2 35	7 39	452 258	537 799	- 2	-	42 89	78 165	- 1	- 2		
Kans.	54	119	1,442	1,826	1	1	583	881	10	3		
S. ATLANTIC	4,846	6,520	31,518	21,165	49	12	31,042	36,557	99	68		
Md.	576	847	2,768	2,402	2	1	420 5,072	4,905	6	- 1		
D.C.	282	431	N	N	- N	-	1,319	60	-	-		
W. Va.	27	50	1,320	4,537 813	N	-	3,118	252	о 5	6		
N.C.	281 270	280 344	6,569	U	13	6	6,098 3 991	7,318	22 17	18 14		
Ga.	683	868	3,377	4,632	15	-	4,453	8,722	Ű	-		
Fla.	2,237	3,228	8,800	8,739	17	-	6,192	6,684	41	24		
E.S. CENTRAL Ky.	609 60	869 152	2,546	2,651	32 10	-	12,436	12,086	146	236		
Ténn.	285	310	4,847	4,818	17	7	4,069	4,271	84	197		
Miss.	113	131	2,902 2,190	487	2	-	2,625	1,070	5 50	26		
W.S. CENTRAL	2,040	2,598	16,494	8,300	13	4	11,188	7,853	122	139		
Ark.	83 385	121 612	474 3 098	620 2 574	1	1	965 2 855	1,539 2 777	- 78	3 58		
Okla.	116	100	2,913	2,673	-	-	1,831	1,709	4	43		
lex.	1,456	1,765	10,009	2,433	9	-	5,537	1,828	40	35		
Mont.	16	785 10	311	5,949 497	36	- 24	2,659	2,978	143	260		
Idaho Wive	18	10	578	615	5	-	43	34	19	67		
Colo.	156	245	101	274	14	8	554	654	18	24		
N. Mex.	58 159	45	1,278	1,564	4 N	3	508	352	26 15	33		
Utah	41	85	660	620	4	-	87	122	2	10		
Nev.	143	155	1,055	1,087	3	3	289	336	3	10		
Wash.	2,973	4,406 309	21,676 3,563	27,698 3,876	69 11	40 4	4,899 780	8,616 900	185 9	164 26		
Oreg.	128	223	1,325	2,094	20	14	205	177	4	3		
Calif. Alaska	2,570	3,784 10	15,675 532	20,666	35	19	3,582 174	7,149 185	109	58 2		
Hawaii	22	80	581	703	Ň	3	158	205	63	75		
Guam PB	2 520	3 122	14 N	162 N	N 21	-	1 249	28	- 10	5 12		
V.I.	29	422	N	N	N	Ŭ	- 240	-	42	-		
Amer. Samoa C.N.M.I.	-	-	N	N	N N	U U	- 11	- 11	2	-		

TABLE II. Provisional cases of selected notifiable diseases, United States,
weeks ending May 17, 1997, and May 18, 1996 (20th Week)

N: Not notifiable U: Unavailable -: no reported cases C.N.M.I.: Commonwealth of Northern Mariana Islands

*Updated monthly to the Division of HIV/AIDS Prevention–Surveillance and Epidemiology, National Center for HIV, STD, and TB Prevention, last update April 29, 1997. *National Electronic Telecommunications System for Surveillance. *Public Health Laboratory Information System.

	Legionellosis		Lyı Dise	me ease	Ma	laria	Syp (Primary &	hilis Secondary)	Tubero	Rabies, Animal	
Reporting Area	Cum. 1997	Cum. 1996	Cum. 1997	Cum. 1996	Cum. 1997	Cum. 1996	Cum. 1997	Cum. 1996	Cum. 1997	Cum. 1996	Cum. 1997
UNITED STATES	308	283	970	1,497	439	423	3,011	4,505	5,184	6,499	2,707
NEW ENGLAND	23	14	193	167	14	13	62	68	127	203	417
Maine N.H.	1 4	1	3 5	23	1	3	-	- 1	2	8 4	87 19
Vt. Mass	3 7	2	2 48	- 14	1 9	1 5	- 35	- 29	2 82	- 57	69 79
R.I.	4	5	33	21	2	3	-	20	7	20	8
MID. ATLANTIC	4 52	62	604	1.175	- 102	- 125	134	203	33 1.125	1.114	581
Upstate N.Y.	11	13	85	559	21	23	14	24	145	118	424
N.J.	6	7	139	134	21	25	49	71	227	255	52
Pa.	35	39 102	375	421	9	9	38	43	156	161	105
Ohio	66	37	17	12	30	53 6	92	295	599 124	107	39 34
Ind. III.	15	25 13	4	4	3 5	3 25	64 24	104 199	51 293	71 400	2 1
Mich. Wis	31 5	18 10	- U	Ū	15 3	10 9	45 47	71 83	88 43	110 29	2
W.N. CENTRAL	28	18	11	36	11	11	51	170	177	163	162
Minn. Iowa	1 5	1	9	1 5	5	3 1	U 3	15 13	47 20	40 19	16 64
Mo.	6	4	-	13	2	5	32	126	71	55	8
N. Dak. S. Dak.	2	2	-	-	-	-	-	-	4 2	2 13	22 17
Nebr. Kans.	9 4	7	2	- 17	1	- 2	- 16	6 10	4 29	12 22	1 34
S. ATLANTIC	45	31	95	51	111	71	1,245	1,468	1,122	1,134	1,160
Del. Md.	4 15	2 5	-72	26 5	2 32	2 20	11 327	16 228	7 109	21 96	25 208
D.C.	2	1	5	1	6	3	41	8	34	51	2
W. Va.	4	9	-	3	-	1	1	192	21	23	240
N.C. S.C.	5 2	3	3 1	10 2	6 5	7 3	273 157	419 173	127 125	125 134	359 57
Ga.	- 13	- 7	1 13	-	12 26	8 19	209 113	282 148	202 386	236 366	108 133
E.S. CENTRAL	9	, 17	25	22	13	13	720	1,110	394	499	106
Ky. Tenn	-	2	2 10	6	2	35	65 302	55 365	77 79	86 165	10 69
Ala.	1	1	2	1	4	1	183	222	163	160	27
WISS.	4	2	4	9	3 5	3 10	374	468 461	75 126	88 753	- 113
Ark.	-	-	-	4	1	-	29	119	76	69	19
La. Okla.	-	2	2	2	4	-	49	206	50	61	46
Tex.	3	-	1	1	-	10	143	74	U	622	47
MOUNTAIN Mont.	16 1	16 1	1	-	28	24	63	52	193	215 7	34 6
ldaho Wyo	1 1	- 2	-	-	- 1	- 2	-	1	4	3	- 11
Colo.	3	5	-	-	14	12	1	16	44	39	-
Ariz.	4	4	- 1	-	4 3	3	54	30	83	30 91	15
Utah Nev.	4 1	1 3	-	-	1 3	3 1	2 6	- 4	6 44	10 34	- 1
PACIFIC	14	20	20	27	125	104	90	221	1,321	1,701	95
Wash. Oreg.	3	1	- 8	1 9	6 8	6 8	6 3	2 4	74 57	94 66	- 1
Calif.	10	19	12	16	107	85	79	214	1,090	1,441	81
Hawaii	- 1	-	-	- 1	2	4	1	- 1	33 67	36 64	-
Guam	-	-	-	-	-	-	-	3	2	45	-
V.I.	-	-	-	-	3 -	-	80 -	48	88 -	56 -	Z I -
Amer. Samoa C.N.M.I.	-	-	-	-	-	-	- 4	- 1	-	-	-

TABLE II. (Cont'd.) Provisional cases of selected notifiable diseases, United States,
weeks ending May 17, 1997, and May 18, 1996 (20th Week)

N: Not notifiable U: Unavailable -: no reported cases

	H. influ	uenzae,	Hepatitis (Viral), by type					Measles (Rubeola)						
	inva	sive		4		В	Indi	genous	lmp	orted [†]	То	Total		
Reporting Area	Cum. 1997*	Cum. 1996	Cum. 1997	Cum. 1996	Cum. 1997	Cum. 1996	1997	Cum. 1997	1997	Cum. 1997	Cum. 1997	Cum. 1996		
UNITED STATES	453	488	9,838	10,204	3,090	3,469	1	31	-	14	45	144		
NEW ENGLAND Maine N.H. Vt	25 3 2	11 - 6	226 35 16	128 10 3	69 4 5	82 2 6	- -	-	-	-	- -	6 - - 1		
Mass.	17	5	97	64	40	19	-	-	-	-	-	4		
R.I. Conn.	2 1	-	20 52	4 44	8 11	4 46	-	-	-	-	-	- 1		
MID. ATLANTIC Upstate N.Y. N.Y. City	52 4 17	96 24 21	749 97 257	693 151 242	416 81 127	576 127 225	- - -	7 1 4	- -	4 3 1	11 4 5	12 4 7		
Pa.	9	29	250	149	103	113	-	1	-	-	1	- 1		
E.N. CENTRAL Ohio Ind.	62 39 5	85 48 3	816 169 123	972 385 136	331 37 35	429 49 53	- - -	4 - -	-	2	6 - -	11 2		
III. Mich. Wis	11 6 1	25 4 5	173 299 52	221 144 86	54 193 12	125 166 36	-	4	-	1 1	5 1	2 2 5		
W.N. CENTRAL	20 12	14 7	747 64	784 35	206	176 13	-	9	-	1	10 1	6		
lowa	2	2	102	166	29	21	-	-	-	-	-	-		
No. N. Dak.	-	- 3	390 7	384 21	136	- 112	-	-	-	-	-	-		
S. Dak. Nebr.	2 1	1 1	12 55	34 93	- 9	- 12	Ū	8	Ū	-	8	-		
Kans.	1	-	117	51	14	18	-	-	-	-	-	-		
S. ATLANTIC Del.	105	90 1	622 11	359 5	455 2	444 2	-	1	-	2	3	2 1		
Md.	36	31 4	116 13	77 15	67 18	67 15	-	-	-	1 1	1 1	-		
Va.	6	4	68	57	44	57	-	-	-	-	-	-		
vv. va. N.C.	3 13	4 14	5 80	43	93	129	-	-	-	-	-	-		
S.C. Ga.	4 17	3 24	53 115	29 13	41 46	38 7	-	-	-	-	-	-		
Fla.	24	5	161	110	138	118	-	1	-	-	1	1		
E.S. CENTRAL Kv.	32 5	17 4	307 27	693 14	276 14	319 32	-	-	2	-	-	-		
Tenn.	19	7	200	495	165	197	-	-	-	-	-	-		
Miss.	-	1	45 35	95	68	20 U	U	-	U	-	-	-		
W.S. CENTRAL	22	17	1,933	1,652	340 17	298	-	3	-	1	4	2		
La.	2	-	82	46	45	39	-	-	-	-	-	-		
Tex.	14	16	662 1,080	733 669	268	208	-	3	-	- 1	4	2		
MOUNTAIN	42	27	1,608	1,586	358	433	1	2	-	-	2	12		
ldaho	- 1	1	44 68	119	5 14	4 53	-	-	-	-	-	1		
Wyo. Colo.	- 6	- 5	18 186	17 149	15 74	13 52	-	-	-	-	-	- 4		
N. Mex.	3	7	109	205	124	142	-	-	-	-	- 2	-		
Utah	3	9 5	288	376	38	50 51	-	-	-	-	-	-		
Nev.	16	-	105	140	17	20	-	-	-	-	-	4		
Wash.	93	131	2,830	3,337 217	20	46	-	5	-	4	9	93 24		
Oreg. Calif.	18 70	18 108	149 2.413	479 2.577	49 553	47 616	-	2	-	- 4	- 6	3 1		
Alaska Hawaii	1	2	16	28	12	1	-	- 2	-	-	- 2	63		
Guam	-	-		30	-	-	- U	-	U	-	-	-		
P.R. V.I.	-	-	139	24	500	74	Ū	-	- U	-	-	1 -		
Amer. Samoa C.N.M.I.	- 4	- 10	- 1	- 1	- 19	- 5	U U	- 1	U U	-	- 1	-		

TABLE III. Provisional cases of selected notifiable diseases preventable by vaccination,
United States, weeks ending May 17, 1997,
and May 18, 1996 (20th Week)

N: Not notifiable U: Unavailable -: no reported cases

 * Of 93 cases among children aged <5 years, serotype was reported for 46 and of those, 18 were type b.

[†]For imported measles, cases include only those resulting from importation from other countries.

Vol. 46 / No. 20

MMWR

Legnoring Jave Cum.		Mening Dis	jococcal ease	Mumps				Pertussis		Rubella			
UNITED STATES 1,590 1,472 30 255 272 114 1,341 1,246 1 20 80 NEW ENGLAND 97 56 - 7 - 8 438 229 - - 11 Mate 9 1 - - - 4 153 17 - - 2 VL 2 3 - - - 4 153 17 - - 2 Mess. 53 19 - 2 - 3 - - 7 R. 6 5 - 4 38 4 138 66 - 1	Reporting Area	Cum. 1997	Cum. 1996	1997	Cum. 1997	Cum. 1996	1997	Cum. 1997	Cum. 1996	1997	Cum. 1997	Cum. 1996	
NEW BOLAND 97 56 - 7 - 8 436 229 - - 1 N.H. 9 1 - - - 1 57 16 - - - - 2 Mis. 56 19 - - - 4 153 7 - - 2 Mis. 56 19 - 2 - 3 2021 133 - - 2 Conn. 18 142 - 24 38 4 138 88 - 1 5 MDATLANTC 134 142 - 207 77 4 160 1283 - - 1 1 N.J. 30 31 - 1 2 26 2 22 160 2 - - - - - - - 1 - 16 160 3 - - - - - - - - - -	UNITED STATES	1,598	1,472	30	255	272	114	1,941	1,246	1	20	80	
Maine 9 8 - - - - 6 10 - Conto 3 3 - - 10 13 1 - 16 4 2 2 2 2 1 - 1 1 1 1 1 1 1 1 1 </td <td>NEW ENGLAND</td> <td>97</td> <td>56</td> <td>-</td> <td>7</td> <td>-</td> <td>8</td> <td>436</td> <td>229</td> <td>-</td> <td>-</td> <td>11</td>	NEW ENGLAND	97	56	-	7	-	8	436	229	-	-	11	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Maine N H	9	8	-	-	-	- 1	6 57	10 16	-	-	-	
Mass. 53 19 - 2 - 3 203 19 - - 7 Can. 18 20 - 1 - 12 3 - - 7 Con. 18 20 - 1 - 18 88 - 1 5 MD.ATEN/TIC 13 23 24 - 0 - 19 14 - 1 1 NJ. 30 31 - - 20 5 4 - 1 1 Pa. 47 51 - 20 77 12 26 2 62 5 4 - 1 Pa. 47 51 - 20 77 72 4 166 183 - 2 <th2< th=""> 2 <th2< th=""> <th2< th=""></th2<></th2<></th2<>	Vt.	2	3	-	-	-	4	153	7	-	-	2	
Conn. 18 20 - 1 - - 5 3 - - 2 Upstare NYR 34 142 - 24 38 4 138 88 - 1 5 NJ, Tory 33 31 - - 2 - 5 4 - - 10 Pa. 47 51 - 20 17 4 64 25 - - - 1 E.N.CENTRAL 218 227 2 27 72 4 160 183 - - - - - 1 - 16 10 18 22 - 2 -	Mass. B I	53	19 5	-	2	-	3	203 12	193	-	-	7	
MID. ATLANTIC 134 142 - 24 38 4 138 88 - 1 5 N.Y. City 23 24 - - 10 - 19 14 - 1 1 Pa. 47 51 - 20 17 4 160 183 - 2 3 Dold 19 14 - 1 <td>Conn.</td> <td>18</td> <td>20</td> <td>-</td> <td>1</td> <td>-</td> <td>-</td> <td>5</td> <td>3</td> <td>-</td> <td>-</td> <td>2</td>	Conn.	18	20	-	1	-	-	5	3	-	-	2	
	MID. ATLANTIC	134	142	-	24	38	4	138	88	-	1	5	
N.J. Pa.30312-541E.N.CENTRAL Dhio218227227724160163-23Ind. 102431-4522212Ind. 112431-45222121Mich. 181828-426-28122-2W.N. CENTRAL 10116108-84-10654 </td <td>N.Y. City</td> <td>34 23</td> <td>36 24</td> <td>-</td> <td>4</td> <td>9 10</td> <td>-</td> <td>50 19</td> <td>45 14</td> <td>-</td> <td>- 1</td> <td>3 1</td>	N.Y. City	34 23	36 24	-	4	9 10	-	50 19	45 14	-	- 1	3 1	
ra. 47 51 - 20 17 4 64 64 65 - - - Dhio 91 75 2 12 26 2 65 -	N.J.	30	31	-	-	2	-	5	4	-	-	1	
	Fa.	4/ 219	10 227	-	20	17	4	04 150	20 192	-	-	-	
	Ohio	91	75	2	12	26	2	62	56	-	-	-	
Mine, 18 29 - 4 26 - 26 12 - - 2 2 WM. CENTRAL 116 108 - 8 4 - 106 54 -	Ind.	24 67	31 67	-	4	5 14	2	22	12 51	-	-	- 1	
Wis. 18 26 - - 1 - 18 52 - 2 - Min. 12 9 - 3 1 - 66 35 - - - Mo. 61 49 - - 1 - 16 10 -	Mich.	18	28	-	4	26	-	26	12	-	-	2	
	Wis.	18	26	-	-	1	-	18	52	-	2	-	
$\begin{array}{c c c c c c c c c c c c c c c c c c c $	W.N. CENTRAL Minn.	116 12	108 9	-	8	4 1	-	106 65	54 35	-	-	-	
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	lowa	24	22	-	3	-	-	15	2	-	-	-	
S. Dak. 3 3 - - - - 1 1 - - - Kans. 10 13 - - - 5 4 - - - S. ATLANTIC 289 220 - 39 29 5 179 112 - 2 1 2 S. ATLANTIC 289 220 - 4 15 3 67 46 - - - 1 1 - - - - 1 1 2 2 1 4 3 - - - 3 2 - - - - 1 1 - - - - 1 1 - - - 1 1 - - - 1 1 0 1 3 - 1 1 1 - - 1 1 1 - - 1 1 1 1 1 1 1 1 1 1 1 <td>N. Dak.</td> <td>1</td> <td>49 2</td> <td>-</td> <td>-</td> <td>2</td> <td>-</td> <td>2</td> <td>-</td> <td>-</td> <td>-</td> <td>-</td>	N. Dak.	1	49 2	-	-	2	-	2	-	-	-	-	
Neb. 3 10 0 2 - 0 2 2 0 - 10	S. Dak.	3	3	-	- 2	-	-	1	1		-	-	
S. ATLANTIC 289 220 - 39 29 5 179 112 - 2 12 Del. 4 2 - 112 - - 1 - <t< td=""><td>Kans.</td><td>10</td><td>13</td><td>-</td><td>-</td><td>-</td><td>-</td><td>2 5</td><td>2 4</td><td>-</td><td>-</td><td>-</td></t<>	Kans.	10	13	-	-	-	-	2 5	2 4	-	-	-	
Del. 4 2 - - - - - 10 - 1 - - - - - 10 - - - - - 10 - - - 10 - - - 10 - - - 10 - - - 10 - - - 10 - - - 10 - - - 10 - - - 10 - - - 10	S. ATLANTIC	289	220	-	39	29	5	179	112	-	2	12	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Del. Md	4 29	2	-	-	- 15	- 3	- 67	10 46	-	-	-	
Va. 25 27 - 4 3 - 19 5 - 1 - N.C. 48 34 - 6 - - 35 24 -	D.C.	1	3	-	-	-	-	2	-	-	-	1	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Va. W. Va.	25 10	27 10	-	4	3	-	19 3	5 2	-	1	-	
S.C., 37 , 31 , -1 , 9 , 4 , $-$, 8 , 1 , $-$, 1 , 1 , $-$, $-$, $-$, $-$, $-$, $-$, $-$, $-$	N.C.	48	34	-	6	-	-	35	24	-	-	-	
Fla.7823-1252391810E.S. CENTRAL122117-1510-36122Tenn.4635-41-167Ala.3033-53-102Miss.1632U46U83U-NWS.CENTRAL1631704282223139-37Ark.222252LaOkla.18145410Okla.18145410MOUNTAIN99892334138159614612441430 <td>S.C. Ga.</td> <td>37 57</td> <td>68</td> <td>-</td> <td>9 4</td> <td>4</td> <td>-</td> <td>8 6</td> <td>6</td> <td>-</td> <td>-</td> <td>-</td>	S.C. Ga.	37 57	68	-	9 4	4	-	8 6	6	-	-	-	
E.S. CENTRAL122117-1510-36122Tenn.4635-41-167Ala.3033-53-102Ala.3033-53-102Miss.1632U46U83U-NW.S. CENTRAL1631704282223139-37Ark.22222521La.2832-78-731Okla.181454Tex.9510242114-1430-366MOUNTAIN998923341381596146124Mont.733Colo.3014-31410225Ariz.222522221-910121 <td>Fla.</td> <td>78</td> <td>23</td> <td>-</td> <td>12</td> <td>5</td> <td>2</td> <td>39</td> <td>18</td> <td>-</td> <td>-</td> <td>10</td>	Fla.	78	23	-	12	5	2	39	18	-	-	10	
ry.30172111 <td>E.S. CENTRAL</td> <td>122</td> <td>117 17</td> <td>-</td> <td>15</td> <td>10</td> <td>-</td> <td>36</td> <td>122</td> <td>-</td> <td>-</td> <td>-</td>	E.S. CENTRAL	122	117 17	-	15	10	-	36	122	-	-	-	
Ala.3033-53-102Miss.1632U46U83U-NWS. CENTRAL1631704282223139-37Ark.22222521La.2832-78-731Okla.181454Tex.9510242114-1430-36MOUNTAIN998923341381596146124Mont.7313Idaho612-2-72450562Vyo133N. Mex.1618NNN22226Nev.68-29-2211 <td< td=""><td>Tenn.</td><td>46</td><td>35</td><td>-</td><td>4</td><td>1</td><td>-</td><td>16</td><td>7</td><td>-</td><td>-</td><td>-</td></td<>	Tenn.	46	35	-	4	1	-	16	7	-	-	-	
N.N.C.N. <th< td=""><td>Ala. Miss.</td><td>30 16</td><td>33 32</td><td>Ū</td><td>5 4</td><td>3</td><td>Ū</td><td>10 8</td><td>2</td><td>Ū</td><td>-</td><td>N</td></th<>	Ala. Miss.	30 16	33 32	Ū	5 4	3	Ū	10 8	2	Ū	-	N	
Ark.2222252La.2832-78-731Okla.18141430-36MOUNTAIN998923341381596146124Mont.73355Idaho612-2-72450562Wyo13Colo.3014-31410225Ariz.222522221-9101211Utah129142-331-Nev.68-29-221-111121Vah129158-137120531Ariz.2229158-137120531Ariz.2229158-137120531N	W.S. CENTRAL	163	170	4	28	22	2	31	39	-	3	7	
La.2832-///31Tex.9510242114-1430-36MOUNTAIN998923341381596146124Mont.73355Idaho612-2-72450562Wyo13Colo.3014-31410225Ariz.222522221-9101211Utah129142-33Nev.68-29-22111111138410269273-103838Wash.4345-58-137120	Ark.	22	22	-		-	2	5	2	-	-	-	
Tex.9510242114-1430-36MOUNTAIN998923341381596146124Mont.73355Idaho612-2-72450562Wyo13Colo.3014-31410225N.Mex.1618NNN22226Ariz.222522221-910121Utah129142-33Nev.68-29-221-1138Wash.4345-58-1371038Oreg.7863-1-41525Calif.2372291566161111118-531Alaska14-22-1Hawaii12-913- <t< td=""><td>La. Okla.</td><td>28 18</td><td>32 14</td><td>-</td><td>-</td><td>8</td><td>-</td><td>7 5</td><td>3</td><td>-</td><td>-</td><td>-</td></t<>	La. Okla.	28 18	32 14	-	-	8	-	7 5	3	-	-	-	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	Tex.	95	102	4	21	14	-	14	30	-	3	6	
Mont.jj <td>MOUNTAIN Mont</td> <td>99 7</td> <td>89</td> <td>23</td> <td>34</td> <td>13</td> <td>81</td> <td>596 5</td> <td>146</td> <td>1</td> <td>2</td> <td>4</td>	MOUNTAIN Mont	99 7	89	23	34	13	81	596 5	146	1	2	4	
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	Idaho	6	12	-	2	-	72	450	56	-	-	2	
N. Mex. 16 18 N N N 1 102 10 1 2 1 Ariz. 22 22 25 22 22 22 1 - 9 10 1 2 1 Utah 12 9 1 4 2 - 3 3 - - - Nev. 6 8 - 2 9 - 2 21 - - 1 PACIFIC 360 343 1 73 84 10 269 273 - 10 38 Wash. 43 45 - 5 8 - 137 120 - - 5 Oreg. 78 63 - 1 - 4 15 25 - - - Alaska 1 4 - 2 2 - 1 - - - - - - - - - - - - -	Wyo. Colo	- 30	- 14	-	1	- 1	- 4	3 102	- 25	-	-	-	
Ariz. 22 22 25 22 22 1 $ 9$ 10 1 2 1 Utah 12 9 1 4 2 $ 3$ 3 $ -$ Nev. 6 8 $ 2$ 9 $ 2$ 21 $ 1$ PACIFIC 360 343 1 73 84 10 269 273 $ 10$ 38 Wash. 43 45 $ 5$ 8 $ 137$ 120 $ 5$ Oreg. 78 63 $ 1$ $ 4$ 15 25 $ -$ Calif. 237 229 1 56 61 6 111 118 $ 5$ 31 Alaska 1 4 $ 2$ 2 $ 1$ $ -$ Hawaii 1 2 $ 9$ 13 $ 5$ 10 $ 5$ 2 Guam $ 1$ U 1 4 U $ U$ $ -$ P.R. 8 2 $ 4$ 1 $ U$ $ -$ V.I. $ U$ $ U$ $ -$ Amer. Samoa $ U$ $ U$ $ U$ $ -$ <	N. Mex.	16	18	N	Ň	Ň	2	22	26	-	-	-	
Nev.68-29-2211PACIFIC3603431738410269273-1038Wash.4345-58-1371205Oreg.7863-1-41525Calif.237229156616111118-531Alaska14-22-1Hawaii12-913-510-52Guam-1U14UUP.R.82-41UV.IUUUAmer. SamoaU1U	Arız. Utah	22 12	25 9	22	22 4	1	-	9 3	10 3	1	2	1	
PACIFIC 360 343 1 73 84 10 269 273 - 10 38 Wash. 43 45 - 5 8 - 137 120 - - 5 Oreg. 78 63 - 1 - 4 15 25 - - - - Calif. 237 229 1 56 61 6 111 118 - 5 31 Alaska 1 4 - 2 2 - 1 -	Nev.	6	8	-	2	9	-	2	21	-	-	1	
Washi. 43 43 - 5 6 - 137 120 - - 5 Oreg. 78 63 - 1 - 4 15 25 - - - - - - - Calif. 237 229 1 56 61 6 111 118 - 5 31 Alaska 1 4 - 2 2 - 1 - <t< td=""><td>PACIFIC</td><td>360</td><td>343</td><td>1</td><td>73</td><td>84</td><td>10</td><td>269</td><td>273</td><td>-</td><td>10</td><td>38</td></t<>	PACIFIC	360	343	1	73	84	10	269	273	-	10	38	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	Oreg.	43 78	45 63	-	5	0 -	4	15	25	-	-	-	
Hawaii 1 2 2 2 1 1 1 1 Hawaii 1 2 - 9 13 - 5 10 - 5 2 Guam - 1 U 1 4 U - - U - <td>Calif. Alaska</td> <td>237 1</td> <td>229 4</td> <td>1</td> <td>56 2</td> <td>61 2</td> <td>6</td> <td>111 1</td> <td>118</td> <td>-</td> <td>5</td> <td>31</td>	Calif. Alaska	237 1	229 4	1	56 2	61 2	6	111 1	118	-	5	31	
Guam - 1 U 1 4 U - - U -	Hawaii	1	2	-	9	13	-	5	10	-	5	2	
HK. 8 2 - 4 1 - - - - VI. - - U - - U - - - Amer. Samoa - - U - - U - - - C.N.M.L - - U 1 - U - - -	Guam	-	1	U	1	4	U	-	-	U	-	-	
Amer. Samoa Ū Ū Ū Ū Ū Ū Ū	P.R. V.I.	8	2	- U	4	1	- U	-	-	- U	-	-	
	Amer. Samoa	-	-	U	- 1	-	U	-	-	U	-	-	

TABLE III. (Cont'd.) Provisional cases of selected notifiable diseases preventable
by vaccination, United States, weeks ending May 17, 1997,
and May 18, 1996 (20th Week)

N: Not notifiable U: Unavailable -: no reported cases

	A	All Causes, By Age (Years)					P&I [†]		All Causes, By Age (Years)						P&I [†]
Reporting Area	All Ages	>65	45-64	25-44	1-24	<1	Total	Reporting Area	All Ages	>65	45-64	25-44	1-24	<1	Total
NEW ENGLAND Boston, Mass. Bridgeport, Conn. Cambridge, Mass. Fall River, Mass. Hartford, Conn. Lowell, Mass. Lynn, Mass. New Bedford, Mass. New Haven, Conn. Providence, R.I. Somerville, Mass. Springfield, Mass.	575 141 38 22 25 67 31 14 29 65 39 65 3 9 U	411 99 25 18 24 41 26 14 22 21 47 3 25 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	106 27 6 2 18 5 - 6 11 7 7 12	32 5 4 2 1 2 - 1 5 7 - 1 U	14 5 3 - 3 - 2 - 1	12 5 	39 9 4 2 2 2 3 1 2 1 4 - 3 U	S. ATLANTIC Atlanta, Ga. Baltimore, Md. Charlotte, N.C. Jacksonville, Fla. Miami, Fla. Norfolk, Va. Richmond, Va. Savannah, Ga. St. Petersburg, Fla. Tampa, Fla. Washington, D.C. Wilmington, Del.	1,302 146 240 97 131 104 57 68 60 55 177 155 12	848 88 148 69 96 60 37 48 30 48 130 87 7	264 35 14 24 24 12 13 22 5 32 38	121 13 31 9 8 18 5 3 1 7 18 5 5	38 5 8 2 1 3 3 3 5 7 -	29 5 8 3 2 1 2 1 2 4 -	64 5 17 6 2 1 2 4 3 2 16 6
Worcester, Mass. MID. ATLANTIC Albany, N.Y. Allentown, Pa. Buffalo, N.Y. Camden, N.J. Elizabeth, N.J. Erie, Pa.	62 2,220 46 58 34 27 53	46 1,550 38 24 44 17 21 41	12 426 5 10 11 3 8	4 174 2 - 2 1 3 2	42 - - 1 2 - 2	27 1 - 1 3 -	6 101 5 1 3 1 3	E.S. CENTRAL Birmingham, Ala. Chattanooga, Tenn. Knoxville, Tenn. Lexington, Ky. Memphis, Tenn. Mobile, Ala. Montgomery, Ala. Nashville, Tenn.	832 171 70 90 102 140 89 43 127	552 117 51 62 65 90 60 31 76	165 33 15 11 22 30 19 5 30	76 11 3 13 8 12 7 6 16	22 9 - 3 3 3 1 3	17 1 4 5 - 2	59 10 4 11 12 2 3 6
Jersey City, N.J. New York City, N.Y. Newark, N.J. Paterson, N.J. Philadelphia, Pa. Pittsburgh, Pa.§ Reading, Pa. Rochester, N.Y. Schenectady, N.Y. Scranton, Pa. Syracuse, N.Y. Trenton, N.J. Utica, N.Y. Yonkers, N.Y.	55 1,207 42 28 301 45 11 118 26 76 30 19 U	43 818 21 16 193 29 11 95 13 23 64 23 16 U	/ 251 11 60 6 - 18 4 2 10 6 1 U	4 104 7 - 33 6 - 3 1 1 2 1 2 U	22 2 12 1 1 - - - - - - - - - - - -	1 12 1 3 3 - - - - - - - U	41 4 19 3 1 6 1 11 1 1	W.S. CENTRAL Austin, Tex. Baton Rouge, La. Corpus Christi, Tex. Dallas, Tex. El Paso, Tex. Houston, Tex. Houston, Tex. Little Rock, Ark. New Orleans, La. San Antonio, Tex. Shreveport, La. Tulsa, Okla.	1,466 77 55 173 73 123 402 71 75 230 41 94	939 51 34 30 100 51 81 247 44 43 159 32 67	313 12 10 12 38 13 27 100 21 17 39 6 18	137 8 11 8 20 4 7 40 2 10 17 2 8	45 2 10 3 4 11 3 5 1	32 2 5 2 4 3 2 10	87 3 2 3 5 5 29 5 20 4 8
E.N. CENTRAL Akron, Ohio Canton, Ohio Chicago, III. Cincinnati, Ohio Cleveland, Ohio Columbus, Ohio Dayton, Ohio Detroit, Mich. Evansville, Ind. Fort Wayne, Ind. Gary, Ind. Grand Banide, Mich	2,020 43 41 408 86 146 154 113 212 50 73 13 55	1,395 31 33 259 60 93 106 87 124 34 52 8 40	382 9 7 83 15 30 35 18 54 8 12 2 8	133 1 43 3 17 2 7 18 6 2 4	58 - 15 5 3 6 - 10 3 2 1 2	51 3 7 3 3 5 1 6 5 1 - 1	129 5 34 10 4 7 6 5 5 5	MOUNTAIN Albuquerque, N.M. Boise, Idaho Colo. Springs, Colo Denver, Colo. Las Vegas, Nev. Ogden, Utah Phoenix, Ariz. Pueblo, Colo. Salt Lake City, Utah Tucson, Ariz. PACIFIC	908 87 43 53 108 201 16 135 28 104 133 1,965	629 66 37 65 126 13 92 24 74 95 1,360	160 9 3 12 20 53 3 18 4 14 24 354	71 8 3 15 14 11 10 10 10	29 2 1 3 6 9 4 3 43	17 2 5 1 - 4 2 1 44	48 2 6 3 4 7 10 3 8 5 146
Grand Rapids, Mich Indianapolis, Ind. Lansing, Mich. Milwaukee, Wis. Peoria, III. Rockford, III. South Bend, Ind. Toledo, Ohio Youngstown, Ohio	179 179 33 125 38 45 49 79 78	40 125 23 90 30 34 38 63 65	8 31 23 7 6 8 11 11	4 14 5 4 1 2 2 2	2 4 - 3 - 1 1 2 -	1 5 1 5 1 3 - 1 -	8 12 2 9 4 2 3 4 3 3	Berkeley, Calif. Fresno, Calif. Glendale, Calif. Honolulu, Hawaii Long Beach, Calif. Los Angeles, Calif. Pasadena, Calif. Portland, Oreg. Sacramento, Calif.	16 73 31 68 63 560 14 132 216	11 47 28 53 39 375 11 98 152	4 14 3 10 16 106 2 19 44	1 4 2 7 52 1 8 11	-7 -3 -12 -3 50	- 1 - 15 - 4 4	1 5 2 7 29 - 11 25
W.N. CENTRAL Des Moines, Iowa Duluth, Minn. Kansas City, Kans. Kansas City, Mo. Lincoln, Nebr. Minneapolis, Minn. Omaha, Nebr. St. Louis, Mo. St. Paul, Minn. Wichita, Kans.	807 73 30 22 113 54 164 92 128 50 81	575 53 25 8 64 45 119 68 97 37 59	128 10 2 8 19 4 25 17 21 10 12	47 6 1 2 7 3 13 2 7 1 5	23 1 2 4 1 3 3 - 2 3	17 3 2 1 4 2 3 - 2	44 2 1 12 13 6 6 4 1	San Diego, Calif. San Francisco, Calif San Jose, Calif. Santa Cruz, Calif. Seattle, Wash. Spokane, Wash. Tacoma, Wash. TOTAL	162 124 188 28 143 56 91 12,095 [¶]	108 85 120 22 97 45 69 8,259	33 24 35 5 21 7 11 2,298	16 14 23 1 18 1 5 955	2 2 5 2 2 314	3 1 8 - 2 1 4 246	16 16 1 7 5 3 717

TABLE IV. Deaths in 122 U.S. cities,* week ending May 17, 1997 (20th Week)

U: Unavailable -: no reported cases *Mortality data in this table are voluntarily reported from 122 cities in the United States, most of which have populations of 100,000 or more. A death is reported by the place of its occurrence and by the week that the death certificate was filed. Fetal deaths are not included. *Pneumonia and influenza. *Because of changes in reporting methods in this Pennsylvania city, these numbers are partial counts for the current week. Complete counts will be available in 4 to 6 weeks. Total includes unknown ages.

Contributors to the Production of the *MMWR* (Weekly)

Weekly Notifiable Disease Morbidity Data and 122 Cities Mortality Data

Denise Koo, M.D., M.P.H.

State Support Team

Robert Fagan Jill Andrews Karl A. Brendel Siobhan Gilchrist, M.P.H. Harry Holden Gerald Jones Felicia Perry Svati Shah, M.P.H.

CDC Operations Team

Carol M. Knowles Deborah A. Adams Willie J. Anderson Christine R. Burgess Timothy M. Copeland Patsy A. Hall Myra A. Montalbano Angela Trosclair, M.S.

Desktop Publishing and Graphics Support

Morie M. Higgins Peter M. Jenkins

The Morbidity and Mortality Weekly Report (MMWR) Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy on Friday of each week, send an e-mail message to *listserv@listserv.cdc.gov*. The body content should read SUBscribe mmwr-toc. Electronic copy also is available from CDC's World-Wide Web server at http://www.cdc.gov/ or from CDC's file transfer protocol server at ftp.cdc.gov. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone (202) 512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to: Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone (404) 332-4555.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

 Director, Centers for Disease Control and Prevention David Satcher, M.D., Ph.D. Deputy Director, Centers for Disease Control and Prevention Claire V. Broome, M.D. Director, Epidemiology Program Office Stephen B. Thacker, M.D., M.Sc. 	Editor, <i>MMWR</i> Series Richard A. Goodman, M.D., M.P.H. Managing Editor, <i>MMWR</i> (weekly) Karen L. Foster, M.A. Writers-Editors, <i>MMWR</i> (weekly) David C. Johnson Darlene D. Rumph Person Teresa F. Rutledge Caran R. Wilbanks							
☆U.S. Government Printing Office: 1997-532-228/67003 Begion IV								